



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 5, 2026 – 03:57 AM UTC

PDB ID : 9IAU / pdb_00009iau
Title : Non-immunogenic staphylokinase variant SY155
Authors : Kasiarova, L.; Marek, M.
Deposited on : 2025-02-11
Resolution : 2.09 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	136	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 14%, orange 14%, orange 71%, yellow 71%, yellow 85%, green 85%, green 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 71% 14% • 14% </div> </div>
1	B	136	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 1%, orange 1%, orange 63%, yellow 63%, yellow 78%, green 78%, green 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 63% 15% • 21% </div> </div>
1	C	136	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 1%, orange 1%, orange 68%, yellow 68%, yellow 83%, green 83%, green 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 68% 15% •• 14% </div> </div>
1	D	136	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 1%, orange 1%, orange 56%, yellow 56%, yellow 71%, green 71%, green 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 56% 15% • 28% </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3739 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Staphylokinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	117	Total	C	N	O	S	0	0	0
			927	603	148	175	1			
1	B	107	Total	C	N	O	S	0	0	0
			839	548	133	157	1			
1	C	117	Total	C	N	O	S	0	0	0
			927	603	148	175	1			
1	D	98	Total	C	N	O	S	0	2	0
			790	516	123	150	1			

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	35	ALA	LYS	conflict	UNP Q99SU7
A	65	GLN	GLU	conflict	UNP Q99SU7
A	74	ARG	LYS	conflict	UNP Q99SU7
A	82	ALA	ASP	conflict	UNP Q99SU7
A	84	ALA	SER	conflict	UNP Q99SU7
A	90	ALA	THR	conflict	UNP Q99SU7
A	99	ASP	GLU	conflict	UNP Q99SU7
A	101	SER	THR	conflict	UNP Q99SU7
A	108	ALA	GLU	conflict	UNP Q99SU7
A	109	ALA	LYS	conflict	UNP Q99SU7
A	130	THR	LYS	conflict	UNP Q99SU7
A	135	ARG	LYS	conflict	UNP Q99SU7
B	35	ALA	LYS	conflict	UNP Q99SU7
B	65	GLN	GLU	conflict	UNP Q99SU7
B	74	ARG	LYS	conflict	UNP Q99SU7
B	82	ALA	ASP	conflict	UNP Q99SU7
B	84	ALA	SER	conflict	UNP Q99SU7
B	90	ALA	THR	conflict	UNP Q99SU7
B	99	ASP	GLU	conflict	UNP Q99SU7
B	101	SER	THR	conflict	UNP Q99SU7
B	108	ALA	GLU	conflict	UNP Q99SU7

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	109	ALA	LYS	conflict	UNP Q99SU7
B	130	THR	LYS	conflict	UNP Q99SU7
B	135	ARG	LYS	conflict	UNP Q99SU7
C	35	ALA	LYS	conflict	UNP Q99SU7
C	65	GLN	GLU	conflict	UNP Q99SU7
C	74	ARG	LYS	conflict	UNP Q99SU7
C	82	ALA	ASP	conflict	UNP Q99SU7
C	84	ALA	SER	conflict	UNP Q99SU7
C	90	ALA	THR	conflict	UNP Q99SU7
C	99	ASP	GLU	conflict	UNP Q99SU7
C	101	SER	THR	conflict	UNP Q99SU7
C	108	ALA	GLU	conflict	UNP Q99SU7
C	109	ALA	LYS	conflict	UNP Q99SU7
C	130	THR	LYS	conflict	UNP Q99SU7
C	135	ARG	LYS	conflict	UNP Q99SU7
D	35	ALA	LYS	conflict	UNP Q99SU7
D	65	GLN	GLU	conflict	UNP Q99SU7
D	74	ARG	LYS	conflict	UNP Q99SU7
D	82	ALA	ASP	conflict	UNP Q99SU7
D	84	ALA	SER	conflict	UNP Q99SU7
D	90	ALA	THR	conflict	UNP Q99SU7
D	99	ASP	GLU	conflict	UNP Q99SU7
D	101	SER	THR	conflict	UNP Q99SU7
D	108	ALA	GLU	conflict	UNP Q99SU7
D	109	ALA	LYS	conflict	UNP Q99SU7
D	130	THR	LYS	conflict	UNP Q99SU7
D	135	ARG	LYS	conflict	UNP Q99SU7

- Molecule 2 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			10	6	4		
2	B	1	Total	C	O	0	0
			10	6	4		
2	B	1	Total	C	O	0	0
			10	6	4		
2	C	1	Total	C	O	0	0
			10	6	4		
2	D	1	Total	C	O	0	0
			10	6	4		
2	D	1	Total	C	O	0	0
			10	6	4		
2	D	1	Total	C	O	0	0
			10	6	4		

- Molecule 3 is CITRATE ANION (CCD ID: FLC) (formula: $C_6H_5O_7$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			13	6	7		
3	D	1	Total	C	O	0	0
			13	6	7		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	54	Total	O	0	0
			54	54		
4	B	34	Total	O	0	0
			34	34		
4	C	40	Total	O	0	0
			40	40		
4	D	32	Total	O	0	0
			32	32		


- Molecule 1: Staphylokinase




- Chain B:  %



- Chain C: 68% 15% 2% 14%



- Chain D:  56% 15% 28%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	46.01Å 79.48Å 79.88Å 90.00° 105.62° 90.00°	Depositor
Resolution (Å)	44.31 – 2.09 44.31 – 2.09	Depositor EDS
% Data completeness (in resolution range)	94.0 (44.31-2.09) 89.2 (44.31-2.09)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.05 (at 2.08Å)	Xtriage
Refinement program	PHENIX 1.14_3260	Depositor
R, R_{free}	0.233 , 0.274 0.235 , 0.273	Depositor DCC
R_{free} test set	1525 reflections (4.64%)	wwPDB-VP
Wilson B-factor (Å ²)	30.4	Xtriage
Anisotropy	0.329	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 47.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	0.125 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3739	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 25.64 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.0509e-03.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, FLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	1/951 (0.1%)	0.46	0/1293
1	B	0.16	0/860	0.44	0/1172
1	C	0.26	0/951	0.62	3/1293 (0.2%)
1	D	0.14	0/808	0.40	0/1098
All	All	0.27	1/3570 (0.0%)	0.49	3/4856 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	71	THR	C-O	-5.51	1.17	1.23

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	95	ASN	CA-CB-CG	7.56	120.16	112.60
1	C	93	ASP	O-C-N	6.22	128.56	122.09
1	C	93	ASP	CA-C-O	-5.74	114.76	120.90

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	120	ILE	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	927	0	933	14	0
1	B	839	0	843	19	0
1	C	927	0	933	22	0
1	D	790	0	787	15	0
2	B	30	0	42	3	0
2	C	10	0	14	0	0
2	D	30	0	42	4	0
3	B	13	0	5	0	0
3	D	13	0	5	0	0
4	A	54	0	0	0	0
4	B	34	0	0	0	0
4	C	40	0	0	1	0
4	D	32	0	0	0	0
All	All	3739	0	3604	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 62 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:118:GLU:HB2	1:C:94:LYS:HD2	1.51	0.92
1:C:96:LYS:HD3	1:C:96:LYS:H	1.37	0.88
1:A:91:TYR:HB2	1:A:120:ILE:HD12	1.56	0.87
1:D:73:TYR:HB2	2:D:203:PGE:H42	1.67	0.76
1:C:96:LYS:H	1:C:96:LYS:CD	1.98	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	115/136 (85%)	112 (97%)	2 (2%)	1 (1%)	14	10
1	B	103/136 (76%)	98 (95%)	3 (3%)	2 (2%)	6	3
1	C	115/136 (85%)	107 (93%)	5 (4%)	3 (3%)	4	1
1	D	92/136 (68%)	85 (92%)	5 (5%)	2 (2%)	5	2
All	All	425/544 (78%)	402 (95%)	15 (4%)	8 (2%)	6	3

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	120	ILE
1	B	121	LYS
1	C	96	LYS
1	D	105	PRO
1	A	120	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	101/117 (86%)	101 (100%)	0	100	100
1	B	91/117 (78%)	91 (100%)	0	100	100
1	C	101/117 (86%)	99 (98%)	2 (2%)	48	56
1	D	88/117 (75%)	88 (100%)	0	100	100
All	All	381/468 (81%)	379 (100%)	2 (0%)	81	88

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	94	LYS
1	C	96	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	43	HIS
1	B	28	ASN
1	C	126	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PGE	D	201	-	9,9,9	0.53	0	8,8,8	0.39	0
2	PGE	B	204	-	9,9,9	0.53	0	8,8,8	0.40	0
2	PGE	B	202	-	9,9,9	0.53	0	8,8,8	0.39	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	FLC	B	203	-	12,12,12	1.09	0	17,17,17	1.28	1 (5%)
3	FLC	D	202	-	12,12,12	1.09	0	17,17,17	1.25	1 (5%)
2	PGE	D	203	-	9,9,9	0.52	0	8,8,8	0.40	0
2	PGE	C	200	-	9,9,9	0.52	0	8,8,8	0.41	0
2	PGE	D	204	-	9,9,9	0.53	0	8,8,8	0.39	0
2	PGE	B	201	-	9,9,9	0.53	0	8,8,8	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PGE	D	201	-	-	1/7/7/7	-
2	PGE	B	204	-	-	2/7/7/7	-
2	PGE	B	202	-	-	3/7/7/7	-
3	FLC	B	203	-	-	0/16/16/16	-
3	FLC	D	202	-	-	2/16/16/16	-
2	PGE	D	203	-	-	4/7/7/7	-
2	PGE	C	200	-	-	4/7/7/7	-
2	PGE	D	204	-	-	1/7/7/7	-
2	PGE	B	201	-	-	3/7/7/7	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	202	FLC	OB2-CBC-CB	3.58	120.02	113.14
3	B	203	FLC	OB2-CBC-CB	3.43	119.72	113.14

There are no chirality outliers.

5 of 20 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	200	PGE	O2-C3-C4-O3
2	B	201	PGE	O2-C3-C4-O3
2	D	203	PGE	O2-C3-C4-O3
2	C	200	PGE	O3-C5-C6-O4
2	B	204	PGE	O1-C1-C2-O2


There are no ring outliers.


4 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	201	PGE	1	0
2	B	204	PGE	1	0
2	B	202	PGE	2	0
2	D	203	PGE	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	117/136 (86%)	-0.95	0 100 100	19, 34, 65, 97	0
1	B	107/136 (78%)	-0.67	2 (1%) 66 69	20, 41, 96, 116	0
1	C	117/136 (86%)	-0.87	0 100 100	18, 41, 83, 118	0
1	D	98/136 (72%)	-0.82	2 (2%) 65 67	15, 33, 88, 113	2 (2%)
All	All	439/544 (80%)	-0.83	4 (0%) 81 83	15, 37, 83, 118	2 (0%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	101	SER	3.1
1	D	104	PHE	3.0
1	B	120	ILE	2.4
1	B	91	TYR	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	PGE	C	200	10/10	0.97	0.07	46,67,72,73	0
2	PGE	D	203	10/10	0.97	0.07	33,44,51,57	0
2	PGE	B	204	10/10	0.98	0.07	54,57,70,70	0
2	PGE	B	201	10/10	0.98	0.07	46,53,68,69	0
2	PGE	D	201	10/10	0.98	0.07	51,58,62,68	0
2	PGE	B	202	10/10	0.98	0.06	41,49,59,65	0
2	PGE	D	204	10/10	0.98	0.06	38,52,60,62	0
3	FLC	B	203	13/13	0.99	0.03	27,30,35,38	0
3	FLC	D	202	13/13	0.99	0.03	21,37,49,52	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around PGE D 203:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around PGE B 204:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PGE B 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around PGE D 201:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PGE B 202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around PGE D 204:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FLC B 203:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FLC D 202:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.