



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 5, 2026 – 02:58 AM UTC

PDB ID : 9IG7 / pdb_00009ig7
Title : KOD-H4 DNA polymerase mutant in a binary complex with DNA:DNA containing two AtNA nucleotides
Authors : Gutfreund, C.; Betz, K.
Deposited on : 2025-02-19
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	774	<div> <div></div> <div>86%</div> <div>11%</div> <div>.</div> </div>
2	P	13	<div> <div>85%</div> <div>15%</div> </div>
3	T	16	<div> <div>44%</div> <div>56%</div> </div>

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 7003 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	756	Total	C	N	O	S	0	3	0
			6224	4001	1057	1149	17			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	93	GLN	VAL	engineered mutation	UNP D0VWU9
A	114	THR	ILE	engineered mutation	UNP D0VWU9
A	141	ALA	ASP	engineered mutation	UNP D0VWU9
A	143	ALA	GLU	engineered mutation	UNP D0VWU9
A	147	HIS	GLU	engineered mutation	UNP D0VWU9
A	383	LYS	SER	engineered mutation	UNP D0VWU9
A	429	GLY	LYS	engineered mutation	UNP D0VWU9
A	445	LEU	PHE	engineered mutation	UNP D0VWU9
A	485	LEU	ALA	engineered mutation	UNP D0VWU9
A	493	VAL	TYR	engineered mutation	UNP D0VWU9
A	496	HIS	TYR	engineered mutation	UNP D0VWU9
A	497	MET	TYR	engineered mutation	UNP D0VWU9
A	499	PHE	TYR	engineered mutation	UNP D0VWU9
A	500	GLU	ALA	engineered mutation	UNP D0VWU9
A	501	ASN	ARG	engineered mutation	UNP D0VWU9
A	521	LEU	ILE	engineered mutation	UNP D0VWU9
A	584	LYS	GLU	engineered mutation	UNP D0VWU9
A	664	LYS	GLU	engineered mutation	UNP D0VWU9
A	726	ARG	LYS	engineered mutation	UNP D0VWU9
A	735	LYS	ASN	engineered mutation	UNP D0VWU9


- Molecule 2 is a DNA chain called DNA (5'-D(*GP*AP*CP*CP*AP*CP*GP*GP*CP*CP*AP*CP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	P	13	Total	C	N	O	P	0	0	0
			261	124	53	72	12			

- Molecule 3 is a DNA chain called DNA/RNA (5'-D(*AP*AP*C)-R(P*(A6U)P*(A6G))-D(P*TP*GP*GP*CP*CP*GP*TP*GP*GP*TP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	T	16	Total	C	N	O	P	0	0	0
			330	157	60	98	15			

- Molecule 4 is IMIDAZOLE (CCD ID: IMD) (formula: C₃H₅N₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	N	0	0
			5	3	2		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is FORMIC ACID (CCD ID: FMT) (formula: CH_2O_2).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			3	1	2		

- Molecule 7 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 8 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	T	1	Total	C	O	0	0
			4	2	2		

- Molecule 9 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	1	Total	Mg	0	0
			1	1		

- Molecule 10 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	T	1	Total	Mn	0	0
			1	1		


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	119	Total	O	0	0
			119	119		
11	P	6	Total	O	0	0
			6	6		
11	T	15	Total	O	0	0
			15	15		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase



- Molecule 2: DNA (5'-D(*GP*AP*CP*CP*AP*CP*GP*GP*CP*CP*AP*CP*A)-3')



- Molecule 3: DNA/RNA (5'-D(*AP*AP*C)-R(P*(A6U)P*(A6G))-D(P*TP*GP*GP*CP*CP*GP*TP*GP*GP*TP*C)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	111.47Å 153.47Å 69.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.37 – 2.60 43.37 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.8 (43.37-2.60) 99.8 (43.37-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.11 (at 2.61Å)	Xtriage
Refinement program	PHENIX 1.21.2_5419	Depositor
R, R_{free}	0.199 , 0.259 0.199 , 0.258	Depositor DCC
R_{free} test set	1097 reflections (1.62%)	wwPDB-VP
Wilson B-factor (Å ²)	64.7	Xtriage
Anisotropy	0.466	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 37.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7003	wwPDB-VP
Average B, all atoms (Å ²)	73.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IMD, FMT, A6U, MG, EDO, MES, A6G, GOL, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.13	0/6360	0.31	0/8576
2	P	0.18	0/293	0.31	0/449
3	T	0.17	0/318	0.36	0/487
All	All	0.13	0/6971	0.31	0/9512

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6224	0	6274	51	0
2	P	261	0	145	1	0
3	T	330	0	184	5	0
4	A	5	0	5	0	0
5	A	18	0	24	0	0
6	A	3	0	1	0	0
7	A	12	0	13	0	0
8	A	4	0	6	0	0
8	T	4	0	6	0	0
9	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
10	T	1	0	0	0	0
11	A	119	0	0	3	0
11	P	6	0	0	0	0
11	T	15	0	0	1	0
All	All	7003	0	6658	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:753:GLU:OE1	1:A:753:GLU:N	2.19	0.73
1:A:58:ARG:NH1	1:A:92:ASP:OD1	2.25	0.69
1:A:317:LYS:NZ	1:A:321:GLU:OE2	2.26	0.68
1:A:70:LYS:HE2	1:A:81:GLU:HB3	1.80	0.64
1:A:713:ARG:NH1	1:A:728:ASP:OD2	2.30	0.62
1:A:174:LYS:HE2	1:A:299:TRP:CE2	2.36	0.61
1:A:554:GLU:O	1:A:557:LYS:N	2.36	0.59
1:A:525:ILE:HG23	1:A:536:VAL:HG21	1.86	0.56
1:A:552:ASP:OD1	1:A:553:ALA:N	2.38	0.56
1:A:743:ARG:HH22	3:T:10:DC:P	2.29	0.56
1:A:23:ASN:N	1:A:133:GLU:OE2	2.40	0.52
1:A:670:LEU:HD13	1:A:684:LYS:HG2	1.91	0.52
1:A:704:LEU:HD11	1:A:715:ILE:HD13	1.91	0.51
1:A:400:ILE:HG22	1:A:547:THR:HB	1.93	0.51
1:A:25:GLU:OE1	1:A:27:LYS:NZ	2.45	0.49
1:A:51:VAL:HA	1:A:54:ILE:HD12	1.94	0.49
1:A:644:LYS:NZ	1:A:756:ARG:O	2.46	0.49
1:A:153:ALA:HA	1:A:218:TYR:CZ	2.48	0.49
1:A:241:ILE:HD13	1:A:250:VAL:HG22	1.94	0.48
1:A:287:LYS:O	11:A:901:HOH:O	2.19	0.48
1:A:637:GLU:H	1:A:637:GLU:CD	2.22	0.48
1:A:158:LEU:HD22	1:A:299:TRP:CE2	2.48	0.48
1:A:738:LEU:HD13	1:A:752:LYS:HB2	1.96	0.48
1:A:280:GLU:OE2	11:A:901:HOH:O	2.20	0.48
1:A:388:TYR:HE1	1:A:390:LYS:HE3	1.79	0.48
1:A:58:ARG:HE	1:A:59:HIS:CE1	2.32	0.47
1:A:558:LYS:NZ	1:A:562:GLU:OE2	2.48	0.47
1:A:730:GLU:OE2	11:A:902:HOH:O	2.20	0.47
3:T:7:DG:H2'	3:T:8:DG:C8	2.50	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:751:ARG:HB3	1:A:753:GLU:OE1	2.13	0.47
1:A:375:LYS:HE3	1:A:379:ARG:HH22	1.80	0.47
1:A:354[A]:GLU:OE2	1:A:503:ARG:NH1	2.46	0.45
1:A:557:LYS:O	1:A:561:MET:HG2	2.16	0.45
1:A:171:ILE:HG22	1:A:190:MET:HG3	1.98	0.44
1:A:269:ASN:OD1	1:A:613:ARG:NH1	2.50	0.44
1:A:271:PRO:HA	1:A:613:ARG:HB3	2.00	0.44
2:P:6:DC:H2''	2:P:7:DG:N7	2.33	0.44
1:A:99:LYS:HD2	1:A:99:LYS:N	2.32	0.44
1:A:638:LYS:O	1:A:642:ILE:HG12	2.19	0.43
3:T:13:DG:N7	11:T:202:HOH:O	2.37	0.43
1:A:157:ILE:HG13	1:A:222:ARG:HG3	2.00	0.43
1:A:394:ARG:NH2	1:A:743:ARG:O	2.46	0.42
1:A:360:LYS:HD2	1:A:452:LEU:HD22	2.00	0.42
1:A:439:HIS:HE1	1:A:511:GLU:OE2	2.02	0.42
1:A:678:PRO:O	1:A:682:VAL:HG22	2.19	0.42
1:A:388:TYR:CE1	1:A:390:LYS:HE3	2.54	0.41
1:A:419:SER:HB2	1:A:450:PRO:HD3	2.02	0.41
3:T:10:DC:H2''	3:T:11:DG:N7	2.36	0.41
3:T:15:DT:H4'	3:T:16:DC:OP1	2.20	0.41
1:A:686:LEU:HD22	1:A:691:VAL:HG21	2.03	0.41
1:A:460:ARG:HB2	1:A:486:ILE:HG21	2.03	0.40
1:A:610:ILE:HG13	1:A:623:GLN:OE1	2.21	0.40
1:A:432:ASP:CG	1:A:443:LYS:HD2	2.46	0.40
1:A:261:TYR:HB3	1:A:262:PRO:HD3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	757/774 (98%)	731 (97%)	26 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	661/673 (98%)	656 (99%)	5 (1%)	73 88

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	44	ASP
1	A	305	LEU
1	A	388	TYR
1	A	516	TRP
1	A	694	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	59	HIS
1	A	304	ASN
1	A	339	GLN
1	A	382	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	A6U	T	4	10,3	19,22,23	3.65	14 (73%)	23,31,34	2.06	8 (34%)
3	A6G	T	5	2,3	23,26,27	3.09	14 (60%)	29,38,41	2.13	11 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A6U	T	4	10,3	-	1/7/26/27	0/2/2/2
3	A6G	T	5	2,3	-	1/7/26/27	0/3/3/3

All (28) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	T	4	A6U	C2-N1	7.14	1.49	1.38
3	T	4	A6U	C2-N3	7.06	1.50	1.38
3	T	5	A6G	C4-N3	6.77	1.49	1.34
3	T	4	A6U	C6-C5	5.83	1.48	1.35
3	T	5	A6G	C2-N3	5.56	1.46	1.33
3	T	5	A6G	O4'-C4'	5.32	1.53	1.43
3	T	4	A6U	O4'-C4'	4.96	1.53	1.43
3	T	5	A6G	C2-N2	4.89	1.45	1.34
3	T	4	A6U	C4-N3	4.27	1.45	1.38
3	T	5	A6G	C2'-C1'	-3.85	1.45	1.53
3	T	4	A6U	C2'-C1'	-3.62	1.46	1.53
3	T	4	A6U	C5'-C4'	-3.44	1.41	1.51
3	T	5	A6G	C5'-C4'	-3.28	1.41	1.51
3	T	5	A6G	C5-N7	-3.02	1.33	1.39
3	T	4	A6U	C6-N1	3.02	1.45	1.38
3	T	4	A6U	O4-C4	-2.87	1.18	1.24
3	T	5	A6G	C2-N1	2.79	1.44	1.37
3	T	5	A6G	C5-C6	2.49	1.53	1.44
3	T	4	A6U	O2-C2	-2.43	1.18	1.23
3	T	5	A6G	C6-N1	2.43	1.43	1.38
3	T	5	A6G	O6-C6	-2.40	1.19	1.23
3	T	5	A6G	O2'-C2'	2.38	1.48	1.43
3	T	5	A6G	O4'-C6'	2.38	1.47	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	T	4	A6U	C5-C4	2.36	1.48	1.43
3	T	4	A6U	O2'-C2'	2.33	1.48	1.43
3	T	5	A6G	C1'-N9	2.29	1.54	1.48
3	T	4	A6U	C1'-N1	2.25	1.53	1.48
3	T	4	A6U	O4'-C6'	2.22	1.47	1.43

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	T	5	A6G	C5-C4-N3	-5.43	119.74	128.39
3	T	4	A6U	C4-N3-C2	-5.42	119.89	126.61
3	T	5	A6G	C2-N3-C4	4.58	120.20	112.30
3	T	4	A6U	N3-C2-N1	3.88	119.94	114.89
3	T	4	A6U	C5-C4-N3	3.56	119.79	114.80
3	T	5	A6G	N9-C4-N3	3.20	132.35	125.95
3	T	5	A6G	C2-N1-C6	-3.09	119.50	125.11
3	T	4	A6U	O4-C4-C5	-2.98	120.02	125.16
3	T	5	A6G	N9-C8-N7	-2.69	108.42	113.40
3	T	5	A6G	C5-C6-N1	2.67	120.04	113.25
3	T	5	A6G	C6'-C1'-N9	-2.55	108.75	112.22
3	T	5	A6G	O6-C6-C5	-2.51	119.91	126.53
3	T	5	A6G	C6'-O4'-C4'	2.46	115.48	112.19
3	T	4	A6U	O2-C2-N1	-2.32	119.78	122.80
3	T	5	A6G	C2'-C3'-C4'	2.29	114.38	110.23
3	T	4	A6U	C1'-N1-C2	2.12	119.96	117.41
3	T	4	A6U	C6'-C1'-N1	-2.08	109.39	112.22
3	T	4	A6U	C3'-C2'-C1'	2.05	114.84	111.37
3	T	5	A6G	C8-N7-C5	2.03	107.88	104.26

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	T	5	A6G	C3'-C4'-C5'-O5'
3	T	4	A6U	C2'-C1'-N1-C2

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	MES	A	805	-	12,12,12	1.16	1 (8%)	15,16,16	0.70	0
4	IMD	A	801	-	5,5,5	0.31	0	5,5,5	0.85	0
6	FMT	A	803	-	2,2,2	0.63	0	1,1,1	0.25	0
5	GOL	A	804	-	5,5,5	0.34	0	5,5,5	0.40	0
8	EDO	T	101	-	3,3,3	0.26	0	2,2,2	0.34	0
5	GOL	A	802	-	5,5,5	0.34	0	5,5,5	0.44	0
8	EDO	A	806	-	3,3,3	0.25	0	2,2,2	0.32	0
5	GOL	A	807	-	5,5,5	0.28	0	5,5,5	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	MES	A	805	-	-	3/6/14/14	0/1/1/1
4	IMD	A	801	-	-	-	0/1/1/1
5	GOL	A	804	-	-	2/4/4/4	-
8	EDO	T	101	-	-	1/1/1/1	-
5	GOL	A	802	-	-	2/4/4/4	-
8	EDO	A	806	-	-	0/1/1/1	-
5	GOL	A	807	-	-	2/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	805	MES	C8-S	3.00	1.81	1.77

There are no bond angle outliers.

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	802	GOL	C1-C2-C3-O3
5	A	804	GOL	C1-C2-C3-O3
5	A	807	GOL	O1-C1-C2-O2
7	A	805	MES	C8-C7-N4-C3
5	A	807	GOL	O1-C1-C2-C3
5	A	802	GOL	O2-C2-C3-O3
5	A	804	GOL	O2-C2-C3-O3
7	A	805	MES	C8-C7-N4-C5
8	T	101	EDO	O1-C1-C2-O2
7	A	805	MES	N4-C7-C8-S

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	756/774 (97%)	0.00	9 (1%) 76 73	24, 70, 106, 138	3 (0%)
2	P	13/13 (100%)	0.00	0 100 100	69, 92, 116, 117	0
3	T	14/16 (87%)	-0.35	0 100 100	64, 79, 109, 109	0
All	All	783/803 (97%)	-0.00	9 (1%) 78 74	24, 70, 108, 138	3 (0%)

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	389	VAL	4.7
1	A	689	ARG	3.2
1	A	386	GLY	2.9
1	A	388	TYR	2.7
1	A	692	LYS	2.6
1	A	387	GLY	2.4
1	A	59	HIS	2.2
1	A	385[A]	GLU	2.2
1	A	61	THR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	A6G	T	5	24/25	0.94	0.07	68,77,84,95	0
3	A6U	T	4	21/22	0.96	0.07	62,66,72,75	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	GOL	A	802	6/6	0.64	0.22	80,81,81,82	0
8	EDO	T	101	4/4	0.65	0.25	74,76,81,87	0
7	MES	A	805	12/12	0.74	0.29	125,129,143,150	0
5	GOL	A	804	6/6	0.75	0.19	90,91,94,101	0
5	GOL	A	807	6/6	0.75	0.22	74,82,93,96	0
8	EDO	A	806	4/4	0.82	0.15	86,88,90,90	0
6	FMT	A	803	3/3	0.87	0.24	80,80,84,85	0
9	MG	A	808	1/1	0.93	0.21	74,74,74,74	0
4	IMD	A	801	5/5	0.94	0.12	59,60,62,62	0
10	MN	T	102	1/1	0.99	0.03	72,72,72,72	0

6.5 Other polymers [i](#)

There are no such residues in this entry.