



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 21, 2026 – 06:51 AM JST

PDB ID : 9L8C / pdb\_00009l8c  
Title : Structure of human TREX1-DNA complex  
Authors : Zhu, J.; Wang, L.; Lin, C.; Zhou, W.  
Deposited on : 2024-12-27  
Resolution : 2.52 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	7383 (2.54-2.50)
Clashscore	190562	8079 (2.54-2.50)
Ramachandran outliers	187476	7944 (2.54-2.50)
Sidechain outliers	187428	7946 (2.54-2.50)
RSRZ outliers	180081	7387 (2.54-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	243	<div><div>4%</div><div><div></div><div>78%</div><div>8%</div><div>•</div><div>13%</div></div></div>
1	B	243	<div><div>3%</div><div><div></div><div>84%</div><div>6%</div><div>•</div><div>9%</div></div></div>
1	E	243	<div><div>%</div><div><div></div><div>75%</div><div>12%</div><div>•</div><div>12%</div></div></div>
1	F	243	<div><div>2%</div><div><div></div><div>75%</div><div>12%</div><div>•</div><div>12%</div></div></div>
2	C	22	<div><div>18%</div><div><div></div><div>41%</div><div>41%</div><div>18%</div></div></div>
2	D	22	<div><div>14%</div><div><div></div><div>45%</div><div>27%</div><div>27%</div></div></div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	G	22	<div><div></div><div>9%</div><div>36%</div><div>27%</div><div>36%</div></div>
2	H	22	<div><div></div><div>9%</div><div>36%</div><div>27%</div><div>36%</div></div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8070 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Three-prime repair exonuclease 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	211	Total	C	N	O	S	0	0	0
			1604	1026	277	289	12			
1	B	220	Total	C	N	O	S	0	1	0
			1671	1066	290	302	13			
1	E	213	Total	C	N	O	S	0	0	0
			1624	1038	282	292	12			
1	F	215	Total	C	N	O	S	0	0	0
			1638	1048	284	294	12			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP Q9NSU2
B	0	SER	-	expression tag	UNP Q9NSU2
E	0	SER	-	expression tag	UNP Q9NSU2
F	0	SER	-	expression tag	UNP Q9NSU2

- Molecule 2 is a DNA chain called DNA (5'-D(P\*GP\*TP\*TP\*GP\*GP\*CP\*CP\*CP\*TP\*CP\*TP\*TP\*TP\*AP\*GP\*GP\*GP\*CP\*CP\*AP\*TP\*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	18	Total	C	N	O	P	0	0	0
			366	173	64	111	18			
2	D	16	Total	C	N	O	P	0	0	0
			327	154	59	98	16			
2	G	14	Total	C	N	O	P	0	0	0
			286	135	51	86	14			
2	H	14	Total	C	N	O	P	0	0	0
			286	135	51	86	14			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	35	Total 35	O 35	0	0
3	B	75	Total 75	O 75	0	0
3	C	6	Total 6	O 6	0	0
3	D	13	Total 13	O 13	0	0
3	E	60	Total 60	O 60	0	0
3	F	58	Total 58	O 58	0	0
3	G	6	Total 6	O 6	0	0
3	H	15	Total 15	O 15	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Three-prime repair exonuclease 1




- Molecule 1: Three-prime repair exonuclease 1



- Molecule 1: Three-prime repair exonuclease 1





- Molecule 1: Three-prime repair exonuclease 1







- Molecule 2: DNA (5'-D(P\*GP\*TP\*TP\*GP\*GP\*CP\*CP\*CP\*TP\*CP\*TP\*TP\*TP\*AP\*GP\*GP\*GP\*CP\*CP\*AP\*TP\*C)-3')




- Molecule 2: DNA (5'-D(P\*GP\*TP\*TP\*GP\*GP\*CP\*CP\*CP\*TP\*CP\*TP\*TP\*TP\*AP\*GP\*GP\*GP\*CP\*CP\*AP\*TP\*C)-3')



- Molecule 2: DNA (5'-D(P\*GP\*TP\*TP\*GP\*GP\*CP\*CP\*CP\*TP\*CP\*TP\*TP\*TP\*AP\*GP\*GP\*GP\*CP\*CP\*AP\*TP\*C)-3')



- Molecule 2: DNA (5'-D(P\*GP\*TP\*TP\*GP\*GP\*CP\*CP\*CP\*TP\*CP\*TP\*TP\*TP\*AP\*GP\*GP\*GP\*CP\*CP\*AP\*TP\*C)-3')



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	44.31Å 92.79Å 161.51Å 90.00° 91.45° 90.00°	Depositor
Resolution (Å)	44.30 – 2.52 44.30 – 2.52	Depositor EDS
% Data completeness (in resolution range)	99.1 (44.30-2.52) 99.0 (44.30-2.52)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.26 (at 2.51Å)	Xtriage
Refinement program	PHENIX 1.21.1_5127	Depositor
R, $R_{free}$	0.196 , 0.232 0.197 , 0.232	Depositor DCC
$R_{free}$ test set	2001 reflections (4.53%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	35.2	Xtriage
Anisotropy	0.677	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 58.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.097 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	8070	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.62% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.15	0/1645	0.34	0/2244
1	B	0.22	0/1719	0.42	0/2348
1	E	0.17	0/1666	0.34	0/2272
1	F	0.19	0/1682	0.36	0/2296
2	C	0.22	0/407	0.38	0/623
2	D	0.21	0/364	0.41	0/557
2	G	0.23	0/318	0.45	0/486
2	H	0.25	0/318	0.42	0/486
All	All	0.19	0/8119	0.37	0/11312

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1604	0	1610	9	0
1	B	1671	0	1678	10	0
1	E	1624	0	1633	19	0
1	F	1638	0	1647	17	0
2	C	366	0	204	7	0
2	D	327	0	181	5	0
2	G	286	0	159	5	0
2	H	286	0	159	6	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	35	0	0	1	0
3	B	75	0	0	5	0
3	C	6	0	0	1	0
3	D	13	0	0	1	0
3	E	60	0	0	4	0
3	F	58	0	0	8	0
3	G	6	0	0	0	0
3	H	15	0	0	1	0
All	All	8070	0	7271	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:73:PRO:O	3:A:301:HOH:O	1.82	0.95
2:C:6:DC:OP2	3:C:101:HOH:O	1.95	0.83
1:E:65:ASP:OD2	3:E:301:HOH:O	1.96	0.83
1:E:195:HIS:ND1	3:E:304:HOH:O	2.16	0.78
1:E:124:HIS:O	3:E:302:HOH:O	2.00	0.78
1:B:149:GLY:O	3:B:301:HOH:O	2.04	0.76
1:F:32:THR:HG22	1:F:73:PRO:HG3	1.68	0.74
1:E:65:ASP:OD1	3:E:303:HOH:O	2.04	0.73
1:F:65:ASP:OD1	3:F:301:HOH:O	2.12	0.67
1:E:32:THR:HG22	1:E:73:PRO:HG3	1.76	0.66
1:B:198:GLU:OE1	3:B:302:HOH:O	2.13	0.65
1:F:217:ARG:NH2	3:F:306:HOH:O	2.23	0.63
1:F:234:GLY:O	3:F:302:HOH:O	2.15	0.60
1:B:183:TYR:OH	3:B:303:HOH:O	2.16	0.57
2:G:22:DC:H6	2:G:22:DC:H5'	1.70	0.57
2:H:1:DG:N7	3:H:101:HOH:O	2.33	0.56
1:F:208:CYS:HB3	1:F:215:LEU:HD22	1.86	0.56
2:H:7:DC:H42	2:H:16:DG:H1	1.50	0.56
1:F:56:THR:HG23	1:F:57:VAL:HG23	1.88	0.55
1:E:176:SER:O	1:E:185:ARG:NH2	2.39	0.55
1:B:52:GLY:O	1:B:54:PRO:HD3	2.07	0.55
2:C:18:DC:H2''	2:C:19:DC:H5''	1.88	0.54
1:E:159:LEU:HD23	1:E:182:ILE:HD13	1.90	0.54
1:A:41:ARG:NH2	1:A:220:ASP:OD2	2.37	0.54
1:B:68:SER:HB3	3:B:362:HOH:O	2.06	0.54

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:15:DG:H1'	2:D:16:DG:N1	2.23	0.53
2:H:17:DG:H1'	2:H:18:DC:H5'	1.90	0.53
2:D:16:DG:H1'	2:D:17:DG:N7	2.24	0.52
1:E:180:GLY:O	1:E:184:THR:OG1	2.25	0.52
2:D:7:DC:H2''	2:D:8:DC:H3'	1.92	0.52
1:A:67:LEU:HD21	1:A:107:LEU:HD21	1.90	0.51
2:G:4:DG:H2''	2:G:5:DG:C8	2.45	0.51
2:H:1:DG:H2''	2:H:2:DT:H5'	1.93	0.51
2:H:16:DG:H2''	2:H:17:DG:H5'	1.92	0.51
1:F:129:TYR:HB3	3:F:315:HOH:O	2.11	0.51
1:E:159:LEU:HD21	1:E:207:ILE:CG2	2.42	0.50
1:F:174:ARG:CZ	1:F:174:ARG:HB2	2.42	0.49
1:F:230:ARG:NE	3:F:316:HOH:O	2.47	0.47
1:E:159:LEU:HD21	1:E:207:ILE:HG21	1.97	0.47
2:C:4:DG:H2''	2:C:5:DG:C8	2.50	0.46
1:F:100:PHE:HA	1:F:104:LEU:HD23	1.97	0.46
1:F:131:PHE:CZ	1:F:152:CYS:HB2	2.50	0.46
2:H:16:DG:H2''	2:H:17:DG:H8	1.80	0.46
1:B:48:PRO:HG3	3:B:367:HOH:O	2.15	0.46
2:D:15:DG:H1'	2:D:16:DG:C6	2.51	0.45
1:E:114:ARG:HD2	3:F:308:HOH:O	2.15	0.45
1:F:54:PRO:N	1:F:55:PRO:HD3	2.31	0.45
1:E:128:ARG:NH2	2:G:4:DG:H21	2.15	0.45
1:E:208:CYS:HB3	1:E:215:LEU:HD22	2.00	0.44
1:B:54:PRO:N	1:B:55:PRO:CD	2.79	0.44
1:B:100:PHE:HA	1:B:104:LEU:HD23	2.00	0.44
2:D:21:DT:OP1	3:D:101:HOH:O	2.21	0.44
1:A:17:PHE:HB2	1:A:36:LEU:HG	1.99	0.44
1:A:27:SER:O	1:A:75:LYS:NZ	2.50	0.44
1:E:15:ILE:HB	1:E:121:LEU:HD23	2.00	0.44
2:G:16:DG:H2''	2:G:17:DG:C8	2.53	0.44
1:E:159:LEU:HD23	1:E:182:ILE:HG21	2.01	0.43
1:E:113:ARG:NH1	1:F:103:ASN:OD1	2.52	0.43
1:F:230:ARG:NH2	3:F:316:HOH:O	2.51	0.43
1:B:159:LEU:HD13	1:B:159:LEU:HA	1.86	0.43
2:C:3:DT:H2''	2:C:4:DG:N7	2.34	0.42
1:E:39:VAL:HG22	1:E:64:VAL:HG12	2.02	0.42
2:C:19:DC:H2''	2:C:20:DA:C8	2.54	0.42
1:E:131:PHE:CZ	1:E:152:CYS:HB2	2.55	0.42
2:C:16:DG:H2''	2:C:17:DG:C8	2.55	0.42
1:E:24:LEU:HD13	2:G:2:DT:C2	2.55	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:138:LEU:HB3	1:A:143:LEU:O	2.21	0.41
1:A:208:CYS:HB3	1:A:215:LEU:HD22	2.01	0.41
1:A:128:ARG:NH2	2:C:4:DG:H21	2.18	0.41
1:A:39:VAL:HG22	1:A:64:VAL:HG12	2.02	0.41
1:F:131:PHE:HB2	1:F:132:PRO:HD3	2.02	0.41
1:F:137:GLU:HA	1:F:140:MET:HE3	2.03	0.41
1:B:213:GLN:O	1:B:217:ARG:HG3	2.21	0.40
1:F:183:TYR:OH	3:F:303:HOH:O	2.16	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	205/243 (84%)	203 (99%)	2 (1%)	0	100	100
1	B	217/243 (89%)	210 (97%)	6 (3%)	1 (0%)	24	41
1	E	207/243 (85%)	203 (98%)	4 (2%)	0	100	100
1	F	209/243 (86%)	203 (97%)	6 (3%)	0	100	100
All	All	838/972 (86%)	819 (98%)	18 (2%)	1 (0%)	48	67

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	50	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	173/199 (87%)	165 (95%)	8 (5%)	24	45
1	B	182/199 (92%)	177 (97%)	5 (3%)	39	65
1	E	176/199 (88%)	168 (96%)	8 (4%)	24	46
1	F	178/199 (89%)	167 (94%)	11 (6%)	16	32
All	All	709/796 (89%)	677 (96%)	32 (4%)	24	46

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	LEU
1	A	17	PHE
1	A	46	SER
1	A	82	SER
1	A	107	LEU
1	A	109	LEU
1	A	153	VAL
1	A	159	LEU
1	B	49	THR
1	B	50	SER
1	B	83	GLU
1	B	109	LEU
1	B	159	LEU
1	E	102	ASP
1	E	109	LEU
1	E	128	ARG
1	E	140	MET
1	E	144	THR
1	E	164	ARG
1	E	184	THR
1	E	194	SER
1	F	78	SER
1	F	82	SER
1	F	102	ASP
1	F	109	LEU
1	F	148	ASP
1	F	159	LEU
1	F	174	ARG
1	F	175	LYS
1	F	181	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	F	194	SER
1	F	230	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	103	ASN
1	B	12	GLN
1	B	103	ASN
1	E	103	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	211/243 (86%)	0.21	10 (4%) 36 33	27, 42, 69, 112	0
1	B	220/243 (90%)	-0.09	7 (3%) 50 47	22, 32, 68, 108	1 (0%)
1	E	213/243 (87%)	-0.02	3 (1%) 73 71	23, 36, 60, 85	0
1	F	215/243 (88%)	-0.05	6 (2%) 55 51	24, 36, 66, 84	0
2	C	18/22 (81%)	1.01	4 (22%) 2 2	41, 88, 144, 145	0
2	D	16/22 (72%)	0.67	3 (18%) 3 2	29, 58, 144, 145	0
2	G	14/22 (63%)	0.77	2 (14%) 6 5	38, 87, 116, 121	0
2	H	14/22 (63%)	0.54	2 (14%) 6 5	32, 58, 107, 139	0
All	All	921/1060 (86%)	0.06	37 (4%) 42 39	22, 37, 80, 145	1 (0%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	70	CYS	4.0
1	B	53	PRO	3.3
1	B	144	THR	3.1
2	C	10	DC	3.0
1	F	54	PRO	3.0
2	H	7	DC	3.0
1	B	54	PRO	2.9
2	D	8	DC	2.9
1	A	8	PRO	2.8
1	E	49	THR	2.8
1	A	163	GLU	2.7
1	E	174	ARG	2.6
1	F	49	THR	2.6
1	A	48	PRO	2.5
1	A	165	ALA	2.5
1	A	144	THR	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	F	7	PRO	2.5
1	B	166	SER	2.4
1	B	52	GLY	2.4
1	F	55	PRO	2.4
2	D	7	DC	2.4
1	B	234	GLY	2.3
1	A	148	ASP	2.3
2	G	18	DC	2.3
1	E	8	PRO	2.3
2	C	6	DC	2.3
1	F	188	GLY	2.3
1	A	210	TRP	2.3
2	D	15	DG	2.2
2	C	1	DG	2.2
1	A	74	GLY	2.2
1	A	164	ARG	2.2
1	F	163	GLU	2.2
2	H	1	DG	2.1
2	G	7	DC	2.1
1	A	234	GLY	2.1
2	C	8	DC	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.