



wwPDB EM Validation Summary Report ⓘ

Apr 6, 2026 – 02:14 AM UTC

PDB ID : 9M76 / pdb_00009m76
EMDB ID : EMD-63676
Title : UHRF1 bound to a mononucleosome in its pre-active state, with the RING domain bound to the SRA domain.
Authors : Naschberger, A.; Baradaran, R.; Sayed, A.; Fischle, W.
Deposited on : 2025-03-09
Resolution : 4.10 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at
<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY





The reported resolution of this entry is 4.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	6458 (3.60 - 4.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	103	
1	F	103	
2	C	130	
2	G	130	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	D	126	<div><div></div><div>6%</div><div>77%</div><div>23%</div></div>
3	H	126	<div><div></div><div>•</div><div>76%</div><div>24%</div></div>
4	A	136	<div><div></div><div>7%</div><div>79%</div><div>•</div><div>21%</div></div>
4	E	136	<div><div></div><div>6%</div><div>72%</div><div>•</div><div>27%</div></div>
5	I	144	<div><div></div><div>•</div><div>95%</div><div>5%</div></div>
6	J	144	<div><div></div><div>•</div><div>98%</div><div>•</div></div>
7	K	793	<div><div></div><div>35%</div><div>70%</div><div>•</div><div>27%</div></div>

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 31161 atoms, of which 14338 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone H4.

Mol	Chain	Residues	Atoms						AltConf	Trace
1	B	83	Total	C	H	N	O	S	0	0
			1388	422	720	132	113	1		
1	F	86	Total	C	H	N	O	S	0	0
			1437	436	743	140	117	1		

- Molecule 2 is a protein called Histone H2A.

Mol	Chain	Residues	Atoms						AltConf	Trace
2	C	109	Total	C	H	N	O		0	0
			1737	528	898	166	145			
2	G	111	Total	C	H	N	O		0	0
			1758	535	907	168	148			

- Molecule 3 is a protein called Histone H2B 1.1.

Mol	Chain	Residues	Atoms						AltConf	Trace
3	D	97	Total	C	H	N	O	S	0	0
			1564	480	798	142	142	2		
3	H	96	Total	C	H	N	O	S	0	0
			1540	474	785	138	141	2		

- Molecule 4 is a protein called Histone H3.

Mol	Chain	Residues	Atoms						AltConf	Trace
4	E	99	Total	C	H	N	O	S	0	0
			1676	515	859	158	141	3		
4	A	108	Total	C	H	N	O	S	0	0
			1833	558	943	176	153	3		

- Molecule 5 is a DNA chain called DNA (144-MER).

Mol	Chain	Residues	Atoms						AltConf	Trace
5	I	144	Total	C	H	N	O	P	0	0
			4556	1395	1618	535	864	144		

- Molecule 6 is a DNA chain called DNA (144-MER).

Mol	Chain	Residues	Atoms						AltConf	Trace
6	J	144	Total	C	H	N	O	P	0	0
			4583	1404	1616	555	864	144		

- Molecule 7 is a protein called E3 ubiquitin-protein ligase UHRF1.

Mol	Chain	Residues	Atoms						AltConf	Trace
7	K	581	Total	C	H	N	O	S	0	0
			9084	2880	4451	839	879	35		



- Molecule 8 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
8	K	5	Total	Zn	0
			5	5	



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Histone H4





- Molecule 1: Histone H4





- Molecule 2: Histone H2A





- Molecule 2: Histone H2A




- Molecule 3: Histone H2B 1.1





- Molecule 3: Histone H2B 1.1





- Molecule 4: Histone H3




- Molecule 4: Histone H3




- Molecule 5: DNA (144-MER)



- Molecule 6: DNA (144-MER)



- Molecule 7: E3 ubiquitin-protein ligase UHRF1



K187	E285	S354	S459	G610	I684	A754	K187
Y188	G286	E355	L462	L615	R685	Q755	Y188
D189	S287	D356	A463	E619	E686	V756	D189
D190	P288	E357	G464	E630	D687	F757	D190
Y191	M289	W358			K688	S758	Y191
P192	V290		E467	GLU	S689	C759	P192
E193	D291	N365	D468	GLN	N690	P760	E193
N194	N292	D366	D469	GLN	A691	A761	N194
G195	P293	A367	V470	GLU	K692	C762	G195
V196	M294	S368		GLY	L693	R763	V196
V197	R295	E369	F476	GLY	W694	Y764	V197
Q198	R296	L372	T479	PHE	N695	D765	Q198
M199	K297	A373	G480	ALA	E696	L766	M199
N200	S298	G374		SER	V697	G767	N200
S201	G299	E375	R484	ARG	L698	R768	S201
R202	P300	E376	ASP	THR	A699	S769	R202
D203	S301	R376	LEU	GLY	S700	Y770	D203
	C302	L377	SER	LYS	G701	L701	
		R378	GLY	LYS	L702	M772	
R209	K303		ASN	LYS	D703	Q773	R209
T210	H304	K381	LYS	LYS	R704	W774	T210
I211	C305	K382	THR	ARG	P705	N775	I211
I212	K306	K383	ALA	LYS	A706	Q776	I212
K213	D307	ALA	GLU	SER	S707	P777	K213
W214	D308	LYS	GLN	ALA	G708	L778	W214
Q215	V309	MET	SER	GLY	S709	Q779	Q215
D216	N310	ALA	C497	GLY	P710	T780	D216
L217	R311	SER		PRO	F711	N783	L217
E218	L312	ALA	A514	SER	Q712	Q784	E218
V219	C313	THR	P515	ARG	L713	L785	V219
G220	R314	SER	I516	GLY	F714		G220
	V315	SER	Q519	SER			
		GLN	E520	PRO			
P229	H319	ARG	E521	ARG	E720	R793	P229
D230	L320	ASP	G521	ARG	T721		D230
N231	C321	TRP	K525	THR	F722		N231
P232		GLY		SER	Q723		P232
K233	R324	K399	R534	LYS	C724		K233
E234	Q325	G400	Y535	THR	I725		E234
	D326	M401	V536	LYS			
F237	P327	A402	R537	VAL	E729		F237
	D328	C403		GLU	L730		
R253		V404	Y559	PRO			R253
		G405		TYR			
				S675	R733		
L261	M332	K408	V562	L676	T736		L261
G262	C333	E409	K570	T677	A678		G262
D263	D334	C410			T737		D263
D264	E335	T411			V738		D264
S265	C336	I412	Y578	Q680	C739		S265
L266	D337	G423	L579	S681	Q740		L266
N267	M338	I424	E586	S682	H741		N267
	A339		R597	L683			
		1449			K745		
F273	L345	D346	E609		D746		F273
V274	P347	N454			C747		V274
D275	P348	D455			L748		D275
					D749		
K279	L349				R750		K279
I280	S350				S751		I280
E281	S351				F752		E281
R282	V352				R753		R282
P283	P353						P283
G284							G284

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	48320	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	1800	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	0.010	Depositor
Minimum map value	-0.005	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.000	Depositor
Recommended contour level	0.00126	Depositor
Map size (Å)	279.93597, 279.93597, 279.93597	wwPDB
Map dimensions	384, 384, 384	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.7289999, 0.7289999, 0.7289999	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, 5CM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	B	0.10	0/675	0.30	0/903
1	F	0.09	0/702	0.26	0/937
2	C	0.10	0/849	0.25	0/1145
2	G	0.11	0/861	0.27	0/1162
3	D	0.09	0/777	0.27	0/1041
3	H	0.10	0/766	0.27	0/1027
4	A	0.10	0/901	0.27	0/1205
4	E	0.10	0/829	0.26	0/1111
5	I	0.22	0/3269	0.46	0/5037
6	J	0.21	0/3331	0.44	0/5143
7	K	0.12	0/4731	0.31	0/6388
All	All	0.16	0/17691	0.36	0/25099

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	668	720	719	0	0
1	F	694	743	742	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	839	898	897	0	0
2	G	851	907	906	1	0
3	D	766	798	797	0	0
3	H	755	785	784	0	0
4	A	890	943	944	1	0
4	E	817	859	858	1	0
5	I	2938	1618	1618	4	0
6	J	2967	1616	1616	2	0
7	K	4633	4451	4449	13	0
8	K	5	0	0	0	0
All	All	16823	14338	14330	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 20 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:K:302:CYS:O	7:K:306:LYS:N	2.24	0.69
6:J:55:DC:N4	6:J:56:DG:O6	2.40	0.54
2:G:119:LYS:NZ	7:K:321:CYS:SG	2.82	0.53
5:I:-56:DC:N4	5:I:-55:DG:O6	2.44	0.51
4:A:59:GLU:OE1	4:A:59:GLU:N	2.38	0.50

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	81/103 (79%)	79 (98%)	2 (2%)	0	100	100
1	F	84/103 (82%)	82 (98%)	2 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	C	107/130 (82%)	106 (99%)	1 (1%)	0	100	100
2	G	109/130 (84%)	107 (98%)	2 (2%)	0	100	100
3	D	95/126 (75%)	93 (98%)	2 (2%)	0	100	100
3	H	94/126 (75%)	92 (98%)	2 (2%)	0	100	100
4	A	104/136 (76%)	103 (99%)	1 (1%)	0	100	100
4	E	97/136 (71%)	97 (100%)	0	0	100	100
7	K	571/793 (72%)	552 (97%)	19 (3%)	0	100	100
All	All	1342/1783 (75%)	1311 (98%)	31 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	69/79 (87%)	69 (100%)	0	100	100
1	F	71/79 (90%)	71 (100%)	0	100	100
2	C	85/102 (83%)	85 (100%)	0	100	100
2	G	86/102 (84%)	86 (100%)	0	100	100
3	D	83/106 (78%)	83 (100%)	0	100	100
3	H	82/106 (77%)	82 (100%)	0	100	100
4	A	94/111 (85%)	94 (100%)	0	100	100
4	E	86/111 (78%)	86 (100%)	0	100	100
7	K	500/687 (73%)	495 (99%)	5 (1%)	68	76
All	All	1156/1483 (78%)	1151 (100%)	5 (0%)	81	83

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
7	K	134	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
7	K	403	CYS
7	K	408	LYS
7	K	619	GLU
7	K	676	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
7	K	742	ASN
7	K	690	ASN
7	K	304	HIS
3	H	79	HIS
7	K	319	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	5CM	I	-62	5,6	18,21,22	0.16	0	24,30,33	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	5CM	I	-62	5,6	-	4/7/21/22	0/2/2/2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	I	-62	5CM	C2'-C1'-N1-C6
5	I	-62	5CM	C2'-C1'-N1-C2
5	I	-62	5CM	O4'-C1'-N1-C6
5	I	-62	5CM	O4'-C1'-N1-C2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	I	-62	5CM	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-63676. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 192




Y Index: 192




Z Index: 192


6.2.2 Raw map



X Index: 192



Y Index: 192




Z Index: 192


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 159




Y Index: 171




Z Index: 214


6.3.2 Raw map



X Index: 159



Y Index: 203




Z Index: 214


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)

6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.00126. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_63676_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 159 nm³; this corresponds to an approximate mass of 143 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)




*Reported resolution corresponds to spatial frequency of 0.244 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.244 \AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.10	-	-
Author-provided FSC curve	4.00	5.79	4.03
Unmasked-calculated*	4.27	7.72	4.36

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-63676 and PDB model 9M76. Per-residue inclusion information can be found in section [3](#) on page [6](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.00126 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.00126).

























9.4 Atom inclusion [i](#)



At the recommended contour level, 71% of all backbone atoms, 67% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.00126) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.6740	 0.2390
A	 0.7980	 0.3620
B	 0.8390	 0.4150
C	 0.6910	 0.2750
D	 0.7180	 0.2800
E	 0.7190	 0.2520
F	 0.6820	 0.2510
G	 0.8300	 0.3860
H	 0.8030	 0.3550
I	 0.8430	 0.2420
J	 0.8460	 0.2480
K	 0.4300	 0.1200

