



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 4, 2026 – 11:43 PM UTC

PDB ID : 9MCJ / pdb_00009mcj
Title : Crystal structure of SARS-Cov-2 main protease G15S mutant in complex with
Leritrelvir
Authors : Li, W.W.; Li, J.
Deposited on : 2025-03-17
Resolution : 1.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1187 (1.74-1.74)
Clashscore	190562	1207 (1.74-1.74)
Ramachandran outliers	187476	1200 (1.74-1.74)
Sidechain outliers	187428	1200 (1.74-1.74)
RSRZ outliers	180081	1188 (1.74-1.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	299	<div> <div>11%</div> <div> <div></div> <div>90%</div> <div>9%</div> </div> </div>
1	B	299	<div> <div>22%</div> <div> <div></div> <div>90%</div> <div>9%</div> </div> </div>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4932 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 3C-like proteinase nsp5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	297	Total	C	N	O	S	0	0	0
			2265	1430	384	430	21			
1	B	299	Total	C	N	O	S	0	0	0
			2261	1431	380	428	22			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	15	SER	GLY	engineered mutation	UNP P0DTC1
B	15	SER	GLY	engineered mutation	UNP P0DTC1

- Molecule 2 is Leritrelvir bound form (CCD ID: 7ON) (formula: $C_{31}H_{46}F_3N_5O_6$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			45	31	3	5	6		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	F	N	O	0	0
			45	31	3	5	6		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	186	Total	O	0	0
			186	186		
3	B	130	Total	O	0	0
			130	130		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3C-like proteinase nsp5



- Molecule 1: 3C-like proteinase nsp5



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	55.19Å 98.03Å 58.98Å 90.00° 108.54° 90.00°	Depositor
Resolution (Å)	55.92 – 1.74 55.92 – 1.74	Depositor EDS
% Data completeness (in resolution range)	97.0 (55.92-1.74) 97.0 (55.92-1.74)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.78 (at 1.74Å)	Xtriage
Refinement program	PHENIX 1.12_2829, PHENIX 1.12_2829	Depositor
R, R_{free}	0.210 , 0.250 0.214 , 0.249	Depositor DCC
R_{free} test set	2829 reflections (4.64%)	wwPDB-VP
Wilson B-factor (Å ²)	19.9	Xtriage
Anisotropy	0.176	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 36.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4932	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 7ON

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	0/2314	0.57	0/3152
1	B	0.32	0/2311	0.55	1/3151 (0.0%)
All	All	0.33	0/4625	0.56	1/6303 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	255	ALA	N-CA-C	-8.36	99.26	110.55

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2265	0	2186	15	0
1	B	2261	0	2163	17	0
2	A	45	0	0	1	0
2	B	45	0	0	0	0
3	A	186	0	0	0	0
3	B	130	0	0	1	0
All	All	4932	0	4349	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:109:GLY:HA2	1:B:200:ILE:HD13	1.74	0.70
1:B:255:ALA:O	1:B:256:GLN:HB3	1.96	0.66
1:A:109:GLY:HA2	1:A:200:ILE:HD13	1.82	0.61
1:B:233:VAL:HG11	1:B:269:LYS:HG3	1.88	0.56
1:A:86:VAL:HG13	1:A:179:GLY:HA2	1.89	0.55
1:A:27:LEU:HD13	1:A:39:PRO:HD2	1.89	0.53
1:A:227:LEU:HD11	1:A:242:LEU:HB2	1.90	0.52
1:A:66:PHE:CE1	1:A:87:LEU:HD21	2.44	0.52
1:B:127:GLN:HG2	3:B:544:HOH:O	2.09	0.51
1:B:140:PHE:HB3	1:B:144:SER:OG	2.12	0.50
1:B:56:ASP:O	1:B:59:ILE:HG22	2.13	0.48
1:B:190:THR:O	1:B:192:GLN:HG3	2.14	0.47
1:A:6:MET:HE3	1:A:6:MET:HB2	1.55	0.47
1:B:115:LEU:HD11	1:B:122:PRO:HB3	1.97	0.46
1:A:115:LEU:HD11	1:A:122:PRO:HB3	1.98	0.46
1:A:113:SER:O	1:A:149:GLY:HA2	2.16	0.46
1:A:4:ARG:NH1	1:B:127:GLN:O	2.48	0.45
1:B:3:PHE:CZ	1:B:296:VAL:HG22	2.53	0.44
1:B:209:TYR:O	1:B:213:ILE:HG13	2.18	0.44
1:A:145:CYS:SG	2:A:401:7ON:C8	3.05	0.44
1:B:207:TRP:CE2	1:B:288:GLU:HB3	2.53	0.43
1:A:55:GLU:O	1:A:59:ILE:HG12	2.18	0.43
1:B:255:ALA:O	1:B:256:GLN:CB	2.63	0.43
1:A:22:CYS:HB2	1:A:42:VAL:O	2.20	0.42
1:A:95:ASN:HB3	1:A:98:THR:OG1	2.20	0.42
1:B:131:ARG:HG2	1:B:135:THR:O	2.20	0.41
1:A:112:PHE:HZ	1:A:136:ILE:HG21	1.85	0.41
1:B:113:SER:O	1:B:149:GLY:HA2	2.21	0.41
1:A:108:PRO:HB3	1:A:132:PRO:HA	2.02	0.40
1:B:299:GLN:HE21	1:B:299:GLN:HB2	1.62	0.40
1:B:17:MET:HG3	1:B:117:CYS:SG	2.62	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	295/299 (99%)	288 (98%)	7 (2%)	0	100	100
1	B	297/299 (99%)	292 (98%)	4 (1%)	1 (0%)	36	23
All	All	592/598 (99%)	580 (98%)	11 (2%)	1 (0%)	43	29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	256	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	249/259 (96%)	248 (100%)	1 (0%)	84	79
1	B	246/259 (95%)	245 (100%)	1 (0%)	84	79
All	All	495/518 (96%)	493 (100%)	2 (0%)	84	79

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	49	MET
1	B	271	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	19	GLN
1	B	214	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	7ON	A	401	1	49,49,49	6.19	29 (59%)	61,71,71	2.31	17 (27%)
2	7ON	B	401	1	49,49,49	6.36	27 (55%)	61,71,71	1.75	13 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	7ON	A	401	1	-	1/46/93/93	0/5/5/5
2	7ON	B	401	1	-	1/46/93/93	0/5/5/5

All (56) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	7ON	C4-N1	24.09	1.59	1.33
2	B	401	7ON	C4-N1	23.77	1.58	1.33
2	A	401	7ON	C10-N4	-16.30	1.26	1.46
2	B	401	7ON	C10-N4	-16.10	1.27	1.46
2	B	401	7ON	C13-C12	-14.07	1.20	1.54
2	A	401	7ON	C13-C12	-13.79	1.21	1.54
2	B	401	7ON	C2-C3	12.84	1.71	1.53
2	A	401	7ON	C2-C3	12.72	1.71	1.53
2	B	401	7ON	C7-C6	12.18	1.68	1.54
2	B	401	7ON	C2-C1	-9.29	1.28	1.54
2	A	401	7ON	C2-C1	-9.20	1.28	1.54
2	B	401	7ON	C7-C8	7.74	1.68	1.52
2	A	401	7ON	C7-C8	7.34	1.67	1.52
2	A	401	7ON	C30-N5	6.83	1.47	1.34
2	B	401	7ON	C30-N5	6.81	1.47	1.34
2	B	401	7ON	C13-C10	6.81	1.65	1.54
2	A	401	7ON	C13-C10	6.77	1.65	1.54
2	B	401	7ON	C16-C13	6.69	1.71	1.54
2	A	401	7ON	C16-C13	6.64	1.71	1.54
2	B	401	7ON	C3-N1	-6.60	1.32	1.46
2	A	401	7ON	C3-N1	-6.47	1.32	1.46
2	A	401	7ON	C9-N3	6.46	1.47	1.34
2	B	401	7ON	C17-N4	6.25	1.48	1.34
2	B	401	7ON	C8-N2	6.15	1.47	1.34
2	B	401	7ON	C9-N3	6.12	1.47	1.34
2	A	401	7ON	C7-C6	6.08	1.61	1.54
2	A	401	7ON	C8-N2	5.95	1.46	1.34
2	A	401	7ON	C17-N4	5.91	1.47	1.34
2	A	401	7ON	C14-C12	5.66	1.68	1.53
2	A	401	7ON	C25-N2	-5.63	1.35	1.46
2	B	401	7ON	C25-N2	-5.54	1.35	1.46
2	B	401	7ON	C14-C12	5.37	1.67	1.53
2	A	401	7ON	C15-C16	-4.75	1.32	1.51
2	B	401	7ON	C15-C16	-4.62	1.33	1.51
2	A	401	7ON	C11-C12	4.07	1.58	1.53
2	B	401	7ON	C27-C26	-4.04	1.35	1.51
2	A	401	7ON	C27-C26	-3.97	1.35	1.51
2	B	401	7ON	C11-C12	3.92	1.58	1.53
2	A	401	7ON	C1-C4	3.86	1.61	1.52
2	B	401	7ON	C1-C4	3.45	1.60	1.52
2	B	401	7ON	C28-C29	-3.41	1.38	1.51
2	A	401	7ON	C5-C1	3.36	1.61	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	7ON	C28-C29	-3.33	1.38	1.51
2	B	401	7ON	C5-C1	3.28	1.61	1.53
2	A	401	7ON	O6-C7	-2.82	1.36	1.42
2	B	401	7ON	O1-C4	-2.82	1.18	1.23
2	A	401	7ON	C26-C25	2.76	1.59	1.52
2	B	401	7ON	C26-C25	2.60	1.59	1.52
2	B	401	7ON	C28-C27	2.56	1.65	1.48
2	A	401	7ON	C28-C27	2.48	1.65	1.48
2	A	401	7ON	O1-C4	-2.32	1.18	1.23
2	B	401	7ON	O2-C8	-2.27	1.19	1.23
2	A	401	7ON	O4-C17	-2.20	1.18	1.22
2	B	401	7ON	O3-C9	-2.07	1.19	1.23
2	A	401	7ON	O3-C9	-2.07	1.19	1.23
2	A	401	7ON	C29-C25	2.03	1.58	1.52

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	7ON	C7-C6-N3	-9.35	92.56	109.93
2	A	401	7ON	C6-N3-C9	-7.33	110.49	123.25
2	B	401	7ON	C7-C6-N3	-5.96	98.86	109.93
2	B	401	7ON	C6-N3-C9	-4.89	114.75	123.25
2	A	401	7ON	O6-C7-C6	4.49	121.09	108.43
2	A	401	7ON	C2-C1-C4	4.00	108.36	102.87
2	A	401	7ON	O2-C8-C7	-3.73	113.42	119.99
2	B	401	7ON	C5-C1-C4	-3.70	104.80	112.81
2	B	401	7ON	C3-N1-C4	-3.66	106.02	113.79
2	A	401	7ON	O3-C9-N3	-3.53	116.63	122.96
2	B	401	7ON	C11-N4-C10	-3.33	108.06	112.41
2	B	401	7ON	O6-C7-C8	3.32	117.58	110.63
2	A	401	7ON	C3-N1-C4	-3.16	107.08	113.79
2	A	401	7ON	C26-C25-N2	-3.08	105.69	111.98
2	B	401	7ON	O6-C7-C6	-2.93	100.18	108.43
2	A	401	7ON	C22-C21-C20	2.82	117.22	111.42
2	A	401	7ON	C25-N2-C8	2.81	127.81	123.29
2	A	401	7ON	C10-C9-N3	2.76	120.56	115.59
2	B	401	7ON	C29-C25-N2	-2.52	106.84	111.98
2	A	401	7ON	C20-C19-C18	-2.51	108.52	111.85
2	A	401	7ON	C23-C24-C19	-2.45	107.52	111.89
2	A	401	7ON	C11-N4-C10	-2.36	109.34	112.41
2	B	401	7ON	C24-C19-C18	-2.29	108.82	111.85
2	B	401	7ON	C2-C1-C4	2.27	105.99	102.87

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	7ON	F1-C31-C30	-2.27	105.06	111.85
2	A	401	7ON	C29-C25-N2	-2.27	107.35	111.98
2	B	401	7ON	C9-C10-N4	-2.21	106.90	112.01
2	B	401	7ON	C10-C9-N3	2.14	119.44	115.59
2	A	401	7ON	C3-C2-C1	2.11	108.80	105.69
2	B	401	7ON	C13-C10-N4	2.09	106.29	103.21

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	7ON	C5-C6-C7-O6
2	B	401	7ON	C5-C6-C7-O6


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	7ON	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand 7ON A 401





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	297/299 (99%)	0.73	33 (11%)	10 14	12, 21, 38, 55	0
1	B	299/299 (100%)	1.13	66 (22%)	2 3	12, 25, 42, 53	0
All	All	596/598 (99%)	0.93	99 (16%)	4 6	12, 24, 40, 55	0

All (99) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	255	ALA	5.6
1	B	223	PHE	4.8
1	B	73	VAL	4.6
1	B	72	ASN	4.5
1	A	283	GLY	4.5
1	B	212	VAL	4.1
1	A	154	TYR	4.0
1	B	224	THR	3.9
1	B	3	PHE	3.9
1	B	213	ILE	3.8
1	B	218	TRP	3.8
1	A	297	VAL	3.7
1	A	276	MET	3.7
1	A	59	ILE	3.6
1	A	153	ASP	3.6
1	A	155	ASP	3.6
1	A	294	PHE	3.6
1	B	46	SER	3.6
1	B	67	LEU	3.5
1	B	49	MET	3.5
1	B	24	THR	3.4
1	B	94	ALA	3.3
1	B	278	GLY	3.3
1	A	190	THR	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	278	GLY	3.2
1	A	46	SER	3.2
1	A	300	CYS	3.2
1	B	282	LEU	3.1
1	A	280	THR	3.1
1	B	71	GLY	3.0
1	B	64	HIS	3.0
1	B	216	ASP	3.0
1	B	93	THR	3.0
1	B	251	GLY	3.0
1	B	294	PHE	2.9
1	A	299	GLN	2.9
1	A	274	ASN	2.9
1	A	56	ASP	2.8
1	B	215	GLY	2.8
1	A	277	ASN	2.8
1	B	214	ASN	2.8
1	B	254	SER	2.8
1	B	222	ARG	2.8
1	B	220	LEU	2.8
1	B	210	ALA	2.7
1	A	50	LEU	2.7
1	B	258	GLY	2.7
1	B	56	ASP	2.7
1	A	60	ARG	2.7
1	B	297	VAL	2.7
1	A	191	ALA	2.7
1	B	280	THR	2.6
1	B	275	GLY	2.6
1	B	298	ARG	2.6
1	B	256	GLN	2.6
1	B	75	LEU	2.6
1	A	196	THR	2.5
1	B	153	ASP	2.5
1	A	57	LEU	2.5
1	A	279	ARG	2.5
1	B	249	ILE	2.5
1	B	66	PHE	2.5
1	B	78	ILE	2.5
1	B	211	ALA	2.5
1	A	4	ARG	2.5
1	B	74	GLN	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	50	LEU	2.4
1	B	252	PRO	2.4
1	B	259	ILE	2.4
1	B	92	ASP	2.4
1	B	68	VAL	2.4
1	A	24	THR	2.4
1	B	25	THR	2.4
1	B	154	TYR	2.4
1	B	70	ALA	2.3
1	B	196	THR	2.3
1	B	257	THR	2.3
1	B	65	ASN	2.3
1	B	277	ASN	2.3
1	B	23	GLY	2.3
1	B	281	ILE	2.3
1	B	42	VAL	2.3
1	B	299	GLN	2.2
1	A	285	ALA	2.2
1	B	62	SER	2.2
1	A	45	THR	2.2
1	B	22	CYS	2.2
1	A	48	ASP	2.2
1	B	209	TYR	2.2
1	A	224	THR	2.2
1	B	155	ASP	2.2
1	B	300	CYS	2.2
1	A	291	PHE	2.1
1	B	44	CYS	2.1
1	B	301	SER	2.1
1	A	207	TRP	2.1
1	A	223	PHE	2.1
1	B	59	ILE	2.1
1	A	238	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	7ON	B	401	45/45	0.86	0.14	20,26,39,41	0
2	7ON	A	401	45/45	0.89	0.12	19,26,34,39	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 7ON B 401:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.