



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 5, 2026 – 02:38 AM UTC

PDB ID : 9NOA / pdb\_00009noa  
Title : Crystal structure of unliganded Fab MS-1805  
Authors : Jain, M.; Wilson, I.A.  
Deposited on : 2025-03-08  
Resolution : 1.77 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1365 (1.78-1.78)
Clashscore	190562	1395 (1.78-1.78)
Ramachandran outliers	187476	1382 (1.78-1.78)
Sidechain outliers	187428	1382 (1.78-1.78)
RSRZ outliers	180081	1365 (1.78-1.78)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	227	<div> <div>7%</div> <div>83% 12% 5%</div> </div>
1	C	227	<div> <div>4%</div> <div>89% 8% .</div> </div>
2	B	215	<div> <div>%</div> <div>96% . .</div> </div>
2	D	215	<div> <div>%</div> <div>89% 9% .</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6955 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MS-1805 Fab Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	216	Total	C	N	O	S	0	1	0
			1642	1042	282	311	7			
1	C	221	Total	C	N	O	S	0	0	0
			1671	1059	288	317	7			

- Molecule 2 is a protein called MS-1805 Fab Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	212	Total	C	N	O	S	0	0	0
			1635	1025	272	332	6			
2	D	210	Total	C	N	O	S	0	0	0
			1618	1013	270	329	6			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	74	Total	O	0	0
			74	74		
3	B	51	Total	O	0	0
			51	51		
3	C	118	Total	O	0	0
			118	118		
3	D	146	Total	O	0	0
			146	146		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: MS-1805 Fab Heavy Chain




#### • Molecule 1: MS-1805 Fab Heavy Chain



#### • Molecule 2: MS-1805 Fab Light Chain



#### • Molecule 2: MS-1805 Fab Light Chain



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	40.57Å 145.40Å 76.73Å 90.00° 91.44° 90.00°	Depositor
Resolution (Å)	26.38 – 1.77 26.38 – 1.77	Depositor EDS
% Data completeness (in resolution range)	98.1 (26.38-1.77) 96.1 (26.38-1.77)	Depositor EDS
$R_{merge}$	0.19	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.92 (at 1.77Å)	Xtriage
Refinement program	PHENIX (1.21.2_5419: ???)	Depositor
R, $R_{free}$	0.204 , 0.231 0.204 , 0.231	Depositor DCC
$R_{free}$ test set	2010 reflections (2.38%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	11.6	Xtriage
Anisotropy	0.309	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 30.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.039 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	6955	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.72% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.19	0/1686	0.41	0/2294
1	C	0.22	0/1714	0.45	0/2333
2	B	0.21	0/1671	0.45	0/2268
2	D	0.21	0/1653	0.45	0/2244
All	All	0.21	0/6724	0.44	0/9139

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1642	0	1604	17	0
1	C	1671	0	1633	12	0
2	B	1635	0	1576	3	0
2	D	1618	0	1562	12	0
3	A	74	0	0	1	0
3	B	51	0	0	0	0
3	C	118	0	0	2	0
3	D	146	0	0	3	0
All	All	6955	0	6375	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:159:LEU:HD11	1:A:182:VAL:HG21	1.61	0.83
1:A:35:HIS:CD2	1:A:95:VAL:HG12	2.27	0.69
1:C:16:ARG:NE	1:C:16:ARG:HA	2.07	0.69
1:A:123:PRO:HD3	1:A:209:LYS:HD3	1.78	0.65
1:A:40:ALA:HB3	1:A:43:LYS:HE3	1.80	0.63
2:D:185:ASP:HA	2:D:188:LYS:HD3	1.81	0.63
1:C:35:HIS:CD2	1:C:95:VAL:HG12	2.39	0.57
1:A:11:VAL:HG21	1:A:147:PRO:HG3	1.86	0.56
1:A:61:ASP:HA	1:A:64:GLN:HG3	1.88	0.56
2:D:197:THR:HG23	3:D:384:HOH:O	2.07	0.54
2:D:197:THR:HG22	2:D:204:PRO:HG3	1.90	0.53
1:C:138:LEU:HD13	1:C:211:VAL:HB	1.91	0.52
1:A:193:THR:HG23	1:A:210:LYS:HE3	1.91	0.51
1:C:35:HIS:NE2	1:C:95:VAL:HG12	2.27	0.49
2:D:22:THR:HG22	2:D:72:THR:HG22	1.94	0.49
1:C:82:MET:HE1	1:C:109:VAL:HG21	1.95	0.49
1:A:33:GLY:O	1:A:95:VAL:HG13	2.16	0.46
2:D:103:LYS:HE2	2:D:105:GLU:HB3	1.98	0.45
1:C:18:LEU:HB3	1:C:82:MET:HE3	1.98	0.45
2:D:22:THR:HG23	3:D:356:HOH:O	2.16	0.45
1:A:35:HIS:CE1	1:A:50:ILE:HD12	2.52	0.45
1:A:151:THR:HG22	1:A:199:ASN:HB3	1.97	0.45
1:C:1:GLN:HG3	1:C:2:VAL:H	1.81	0.45
1:A:93:VAL:CG1	1:A:100(F):PHE:HB3	2.47	0.45
2:D:145:LYS:HB3	2:D:197:THR:OG1	2.17	0.45
2:B:21:LEU:HD22	2:B:102:THR:HG21	2.00	0.44
2:D:146:VAL:HG23	2:D:194:CYS:SG	2.58	0.44
2:B:120:PRO:HD3	2:B:132:VAL:HG22	2.00	0.44
1:C:108:MET:HE1	3:C:418:HOH:O	2.18	0.44
1:A:35:HIS:HD2	1:A:95:VAL:HG12	1.80	0.43
2:D:147:GLN:HG3	3:D:384:HOH:O	2.17	0.43
1:A:57:LYS:HE2	3:A:304:HOH:O	2.19	0.43
1:A:151:THR:CG2	1:A:199:ASN:HB3	2.49	0.43
2:D:120:PRO:HD3	2:D:132:VAL:HG22	2.01	0.43
1:A:94:ARG:HB3	1:A:102:LEU:HB3	2.01	0.42
2:D:169:LYS:C	2:D:169:LYS:HD3	2.45	0.42
1:A:189:LEU:HD23	1:A:189:LEU:HA	1.92	0.42
1:C:127:SER:HB2	1:C:129:LYS:HD2	2.01	0.42
1:C:87:THR:HG23	1:C:110:THR:HA	2.02	0.41
1:A:119:PRO:HB3	1:A:145:TYR:HB3	2.01	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:13:ALA:HB2	2:D:19:VAL:HG11	2.03	0.41
1:C:119:PRO:HB3	1:C:145:TYR:HB3	2.03	0.41
2:B:142:ARG:NH2	2:B:163:VAL:HG21	2.36	0.40
1:C:95:VAL:HG23	3:C:383:HOH:O	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/227 (93%)	208 (99%)	3 (1%)	0	100	100
1	C	217/227 (96%)	213 (98%)	4 (2%)	0	100	100
2	B	208/215 (97%)	201 (97%)	7 (3%)	0	100	100
2	D	206/215 (96%)	200 (97%)	6 (3%)	0	100	100
All	All	842/884 (95%)	822 (98%)	20 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	182/191 (95%)	179 (98%)	3 (2%)	55	38
1	C	185/191 (97%)	182 (98%)	3 (2%)	55	38

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	187/189 (99%)	187 (100%)	0	100	100
2	D	185/189 (98%)	183 (99%)	2 (1%)	65	51
All	All	739/760 (97%)	731 (99%)	8 (1%)	65	51

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	95	VAL
1	A	109	VAL
1	A	191	THR
1	C	17	SER
1	C	19	ARG
1	C	129	LYS
2	D	12	SER
2	D	56	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	13	GLN
1	A	64	GLN
1	A	199	ASN
1	C	199	ASN
2	D	37	GLN
2	D	152	ASN
2	D	189	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	216/227 (95%)	0.46	16 (7%) 20 24	8, 19, 44, 55	1 (0%)
1	C	221/227 (97%)	0.29	10 (4%) 38 44	7, 16, 32, 52	0
2	B	212/215 (98%)	0.01	3 (1%) 73 80	9, 15, 29, 38	0
2	D	210/215 (97%)	-0.09	3 (1%) 73 80	7, 14, 26, 40	0
All	All	859/884 (97%)	0.17	32 (3%) 45 51	7, 16, 34, 55	1 (0%)

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	213	PRO	5.0
2	D	214	CYS	4.3
1	C	15	GLY	4.2
2	B	95(A)	PHE	3.7
1	C	128	SER	3.5
1	A	191	THR	3.3
1	A	204	ASN	3.1
1	C	100(A)	PRO	3.0
1	A	1	GLN	3.0
1	C	1	GLN	2.9
1	A	188	SER	2.8
1	A	134	GLY	2.8
1	A	127	SER	2.8
1	C	16	ARG	2.8
1	A	100(B)	HIS	2.6
1	C	130	SER	2.6
1	A	187	SER	2.6
1	A	190	GLY	2.5
1	C	129	LYS	2.5
2	D	154	LEU	2.4
1	A	189	LEU	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	100	GLY	2.4
1	C	100(C)	GLY	2.4
1	A	185	PRO	2.3
2	B	127	SER	2.3
1	A	186	SER	2.1
2	D	169	LYS	2.1
1	A	100(C)	GLY	2.1
1	C	133	GLY	2.1
1	A	199	ASN	2.1
1	A	213	PRO	2.0
2	B	214	CYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.