



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 5, 2026 – 03:08 AM UTC

PDB ID : 9PC5 / pdb\_00009pc5  
Title : Crystal structure of Rv0097 with 5 mM CADA soaked (CADA not bound)  
Authors : Ye, N.; Drennan, C.L.  
Deposited on : 2025-06-27  
Resolution : 1.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1187 (1.74-1.74)
Clashscore	190562	1207 (1.74-1.74)
Ramachandran outliers	187476	1200 (1.74-1.74)
Sidechain outliers	187428	1200 (1.74-1.74)
RSRZ outliers	180081	1188 (1.74-1.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	302	
1	B	302	
1	C	302	
1	D	302	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ACT	B	402	-	-	X	-

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 10108 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called (3R)-3-[(carboxymethyl)amino]fatty acid oxygenase/decarboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	288	Total	C	N	O	S	59	5	0
			2321	1487	398	428	8			
1	B	285	Total	C	N	O	S	137	1	0
			2281	1461	394	418	8			
1	C	282	Total	C	N	O	S	24	2	0
			2269	1454	393	414	8			
1	D	289	Total	C	N	O	S	26	2	0
			2312	1481	399	424	8			

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	290	LYS	-	expression tag	UNP P9WG83
A	291	LEU	-	expression tag	UNP P9WG83
A	292	ALA	-	expression tag	UNP P9WG83
A	293	ALA	-	expression tag	UNP P9WG83
A	294	ALA	-	expression tag	UNP P9WG83
A	295	LEU	-	expression tag	UNP P9WG83
A	296	GLU	-	expression tag	UNP P9WG83
A	297	HIS	-	expression tag	UNP P9WG83
A	298	HIS	-	expression tag	UNP P9WG83
A	299	HIS	-	expression tag	UNP P9WG83
A	300	HIS	-	expression tag	UNP P9WG83
A	301	HIS	-	expression tag	UNP P9WG83
A	302	HIS	-	expression tag	UNP P9WG83
B	290	LYS	-	expression tag	UNP P9WG83
B	291	LEU	-	expression tag	UNP P9WG83
B	292	ALA	-	expression tag	UNP P9WG83
B	293	ALA	-	expression tag	UNP P9WG83
B	294	ALA	-	expression tag	UNP P9WG83
B	295	LEU	-	expression tag	UNP P9WG83
B	296	GLU	-	expression tag	UNP P9WG83

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	297	HIS	-	expression tag	UNP P9WG83
B	298	HIS	-	expression tag	UNP P9WG83
B	299	HIS	-	expression tag	UNP P9WG83
B	300	HIS	-	expression tag	UNP P9WG83
B	301	HIS	-	expression tag	UNP P9WG83
B	302	HIS	-	expression tag	UNP P9WG83
C	290	LYS	-	expression tag	UNP P9WG83
C	291	LEU	-	expression tag	UNP P9WG83
C	292	ALA	-	expression tag	UNP P9WG83
C	293	ALA	-	expression tag	UNP P9WG83
C	294	ALA	-	expression tag	UNP P9WG83
C	295	LEU	-	expression tag	UNP P9WG83
C	296	GLU	-	expression tag	UNP P9WG83
C	297	HIS	-	expression tag	UNP P9WG83
C	298	HIS	-	expression tag	UNP P9WG83
C	299	HIS	-	expression tag	UNP P9WG83
C	300	HIS	-	expression tag	UNP P9WG83
C	301	HIS	-	expression tag	UNP P9WG83
C	302	HIS	-	expression tag	UNP P9WG83
D	290	LYS	-	expression tag	UNP P9WG83
D	291	LEU	-	expression tag	UNP P9WG83
D	292	ALA	-	expression tag	UNP P9WG83
D	293	ALA	-	expression tag	UNP P9WG83
D	294	ALA	-	expression tag	UNP P9WG83
D	295	LEU	-	expression tag	UNP P9WG83
D	296	GLU	-	expression tag	UNP P9WG83
D	297	HIS	-	expression tag	UNP P9WG83
D	298	HIS	-	expression tag	UNP P9WG83
D	299	HIS	-	expression tag	UNP P9WG83
D	300	HIS	-	expression tag	UNP P9WG83
D	301	HIS	-	expression tag	UNP P9WG83
D	302	HIS	-	expression tag	UNP P9WG83

- Molecule 2 is ACETATE ION (CCD ID: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is FE (III) ION (CCD ID: FE) (formula: Fe) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Fe	0	0
			1	1		
3	B	1	Total	Fe	0	0
			1	1		
3	C	1	Total	Fe	0	0
			1	1		
3	D	1	Total	Fe	0	0
			1	1		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	C	O	0	0
			7	4	3		
4	D	1	Total	C	O	0	0
			7	4	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	157	Total	O	0	0
			157	157		
5	B	128	Total	O	0	0
			128	128		
5	C	299	Total	O	0	0
			299	299		
5	D	303	Total	O	0	0
			303	303		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: (3R)-3-[(carboxymethyl)amino]fatty acid oxygenase/decarboxylase



- Molecule 1: (3R)-3-[(carboxymethyl)amino]fatty acid oxygenase/decarboxylase



- Molecule 1: (3R)-3-[(carboxymethyl)amino]fatty acid oxygenase/decarboxylase



- Molecule 1: (3R)-3-[(carboxymethyl)amino]fatty acid oxygenase/decarboxylase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	89.54Å 63.78Å 107.94Å 90.00° 98.00° 90.00°	Depositor
Resolution (Å)	29.56 – 1.74 29.56 – 1.74	Depositor EDS
% Data completeness (in resolution range)	96.9 (29.56-1.74) 96.9 (29.56-1.74)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.05 (at 1.74Å)	Xtriage
Refinement program	PHENIX 1.21.2_5419	Depositor
R, $R_{free}$	0.196 , 0.234 0.196 , 0.234	Depositor DCC
$R_{free}$ test set	6000 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.0	Xtriage
Anisotropy	0.430	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 43.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	10108	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 26.16 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.7711e-03.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, ACT, FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.25	0/2404	0.49	0/3280
1	B	0.20	0/2350	0.44	0/3204
1	C	0.30	0/2341	0.52	0/3190
1	D	0.32	0/2385	0.56	0/3252
All	All	0.27	0/9480	0.51	0/12926

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2321	0	2302	4	0
1	B	2281	0	2255	9	0
1	C	2269	0	2248	8	0
1	D	2312	0	2295	6	0
2	A	4	0	3	0	0
2	B	8	0	6	2	0
2	D	8	0	6	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	C	7	0	10	2	0
4	D	7	0	10	0	0
5	A	157	0	0	0	0
5	B	128	0	0	3	0
5	C	299	0	0	1	0
5	D	303	0	0	0	0
All	All	10108	0	9135	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:275:ARG:HH12	2:B:402:ACT:H1	1.50	0.77
1:A:122:ARG:HD3	1:A:271:LEU:HD22	1.78	0.63
1:C:8:GLU:H	4:C:401:PEG:H31	1.63	0.61
1:D:39:LEU:HD11	1:D:251[A]:ILE:HG23	1.82	0.61
1:D:133:TRP:CH2	1:D:141:ARG:HB2	2.36	0.60
1:C:237:SER:HB3	1:C:240:ILE:HG13	1.85	0.57
1:C:146:GLY:HA3	1:C:211:LYS:HE2	1.86	0.56
1:B:251:ILE:HD11	5:B:536:HOH:O	2.10	0.51
1:B:268:ALA:HA	5:B:616:HOH:O	2.11	0.50
1:A:118:PRO:HG2	1:A:122:ARG:NH1	2.25	0.50
1:B:42:LEU:HB2	1:B:250:ILE:HB	1.94	0.49
1:A:95:PHE:C	1:A:261:ARG:HG3	2.38	0.48
2:B:402:ACT:H2	5:B:504:HOH:O	2.13	0.48
1:D:35:TYR:CD2	1:D:192:LYS:HG2	2.49	0.48
1:C:35:TYR:CD2	1:C:192:LYS:HG2	2.48	0.48
1:C:6:LYS:HG2	5:C:711:HOH:O	2.16	0.45
1:B:237:SER:HB3	1:B:240:ILE:HG13	1.99	0.45
1:D:156:HIS:NE2	1:D:203:THR:HG22	2.33	0.43
1:D:133:TRP:CZ3	1:D:141:ARG:HB2	2.54	0.43
1:B:254:ASP:HB3	1:B:257:VAL:HG12	2.01	0.43
1:B:19:PRO:HB2	1:B:51:GLU:OE1	2.19	0.42
1:B:210:ASP:OD2	1:B:214:ASN:HB2	2.20	0.41
1:A:112[B]:VAL:HG23	1:A:251:ILE:HB	2.03	0.41
1:C:122:ARG:HD2	1:C:263:LYS:O	2.20	0.41
1:B:62:ILE:HG21	1:B:77:GLU:HB3	2.03	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:146:GLY:O	1:C:211:LYS:HG2	2.21	0.40
1:C:8:GLU:H	4:C:401:PEG:H21	1.86	0.40
1:D:133:TRP:CH2	1:D:187:VAL:HG21	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	292/302 (97%)	285 (98%)	7 (2%)	0	100	100
1	B	282/302 (93%)	277 (98%)	5 (2%)	0	100	100
1	C	280/302 (93%)	275 (98%)	5 (2%)	0	100	100
1	D	289/302 (96%)	283 (98%)	6 (2%)	0	100	100
All	All	1143/1208 (95%)	1120 (98%)	23 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	256/261 (98%)	256 (100%)	0	100	100
1	B	249/261 (95%)	247 (99%)	2 (1%)	73	64

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	248/261 (95%)	248 (100%)	0	100	100
1	D	253/261 (97%)	252 (100%)	1 (0%)	84	79
All	All	1006/1044 (96%)	1003 (100%)	3 (0%)	86	82

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	203	THR
1	B	211	LYS
1	D	203	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	13	GLN
1	C	236	GLN
1	C	243	GLN
1	D	13	GLN
1	D	244	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 11 ligands modelled in this entry, 4 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	ACT	D	403	3	3,3,3	1.12	0	3,3,3	1.52	1 (33%)
2	ACT	D	402	-	3,3,3	1.21	0	3,3,3	0.90	0
2	ACT	A	401	-	3,3,3	1.16	0	3,3,3	1.07	0
4	PEG	C	401	-	6,6,6	0.26	0	5,5,5	0.27	0
4	PEG	D	401	-	6,6,6	0.27	0	5,5,5	0.26	0
2	ACT	B	401	-	3,3,3	1.11	0	3,3,3	1.16	0
2	ACT	B	402	-	3,3,3	1.22	0	3,3,3	0.95	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	D	401	-	-	0/4/4/4	-
4	PEG	C	401	-	-	1/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	403	ACT	OXT-C-O	2.06	129.68	122.03

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	C	401	PEG	O2-C3-C4-O4

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	401	PEG	2	0
2	B	402	ACT	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	288/302 (95%)	0.34	5 (1%) 69 76	17, 34, 57, 69	20 (6%)
1	B	285/302 (94%)	0.64	10 (3%) 47 56	17, 41, 65, 82	36 (12%)
1	C	282/302 (93%)	-0.06	1 (0%) 88 92	13, 24, 44, 64	9 (3%)
1	D	289/302 (95%)	-0.09	2 (0%) 84 89	12, 24, 40, 56	9 (3%)
All	All	1144/1208 (94%)	0.21	18 (1%) 70 77	12, 31, 56, 82	74 (6%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	145	ARG	3.1
1	B	25	ILE	3.1
1	B	264	HIS	3.0
1	B	267	ALA	2.9
1	B	268	ALA	2.9
1	C	2	THR	2.8
1	A	213	GLY	2.6
1	D	2	THR	2.5
1	B	3	LEU	2.5
1	B	144	ALA	2.3
1	B	216	VAL	2.3
1	D	264	HIS	2.3
1	A	212	ASP	2.3
1	A	268	ALA	2.2
1	B	210	ASP	2.1
1	A	267	ALA	2.1
1	B	19	PRO	2.1
1	B	289	ALA	2.0



## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	PEG	C	401	7/7	0.68	0.14	45,45,50,58	0
2	ACT	B	402	4/4	0.83	0.15	33,34,37,38	0
4	PEG	D	401	7/7	0.84	0.10	32,38,42,46	0
2	ACT	D	403	4/4	0.91	0.14	18,24,26,29	0
2	ACT	D	402	4/4	0.93	0.08	20,24,25,31	0
2	ACT	B	401	4/4	0.93	0.10	28,32,33,33	0
2	ACT	A	401	4/4	0.94	0.12	34,34,37,37	0
3	FE	D	404	1/1	0.99	0.02	17,17,17,17	0
3	FE	A	402	1/1	0.99	0.03	24,24,24,24	0
3	FE	B	403	1/1	0.99	0.03	26,26,26,26	0
3	FE	C	402	1/1	1.00	0.02	17,17,17,17	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around FE D 404:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FE A 402:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around FE B 403:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers ⓘ

There are no such residues in this entry.