



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 5, 2026 – 02:13 AM UTC

PDB ID : 9RSW / pdb_00009rsw
Title : Crystal structure of the human METTL3-METTL14 in complex with compound 4
Authors : Bedi, R.K.; Caflisch, A.
Deposited on : 2025-07-01
Resolution : 2.56 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.56 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1853 (2.58-2.54)
Clashscore	190562	1897 (2.58-2.54)
Ramachandran outliers	187476	1875 (2.58-2.54)
Sidechain outliers	187428	1875 (2.58-2.54)
RSRZ outliers	180081	1853 (2.58-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	228	<div> <div>5%</div> <div> <div></div> <div>70%</div> <div>19%</div> <div>11%</div> </div> </div>
2	B	307	<div> <div>6%</div> <div> <div></div> <div>63%</div> <div>13%</div> <div>•</div> <div>22%</div> </div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3402 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N6-adenosine-methyltransferase catalytic subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	203	Total	C	N	O	S	0	0	0
			1561	1005	270	277	9			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	353	MET	-	initiating methionine	UNP Q86U44

- Molecule 2 is a protein called N(6)-adenosine-methyltransferase non-catalytic subunit METTL14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	238	Total	C	N	O	S	0	0	0
			1794	1152	307	324	11			

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	89	MET	-	initiating methionine	UNP Q9HCE5
B	90	HIS	-	expression tag	UNP Q9HCE5
B	91	HIS	-	expression tag	UNP Q9HCE5
B	92	HIS	-	expression tag	UNP Q9HCE5
B	93	HIS	-	expression tag	UNP Q9HCE5
B	94	HIS	-	expression tag	UNP Q9HCE5
B	95	HIS	-	expression tag	UNP Q9HCE5
B	96	SER	-	expression tag	UNP Q9HCE5
B	97	SER	-	expression tag	UNP Q9HCE5
B	98	GLY	-	expression tag	UNP Q9HCE5
B	99	ARG	-	expression tag	UNP Q9HCE5
B	100	GLU	-	expression tag	UNP Q9HCE5
B	101	ASN	-	expression tag	UNP Q9HCE5
B	102	LEU	-	expression tag	UNP Q9HCE5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	103	TYR	-	expression tag	UNP Q9HCE5
B	104	PHE	-	expression tag	UNP Q9HCE5
B	105	GLN	-	expression tag	UNP Q9HCE5
B	106	GLY	-	expression tag	UNP Q9HCE5

- Molecule 3 is 8-[4-[(4,4-dimethylpiperidin-1-yl)methyl]-2-oxidanyl-phenyl]-2-[6-(methylanino)pyrimidin-4-yl]-2,5,8-triazaspiro[3.5]nonan-6-one (CCD ID: A1JJH) (formula: C₂₅H₃₅N₇O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			34	25	7	2		

- Molecule 4 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Mg	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	3	Total	O	0	0
			3	3		
6	B	5	Total	O	0	0
			5	5		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: N6-adenosine-methyltransferase catalytic subunit



- Molecule 2: N(6)-adenosine-methyltransferase non-catalytic subunit METTL14



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	64.19Å 64.19Å 227.34Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	44.82 – 2.56 44.82 – 2.56	Depositor EDS
% Data completeness (in resolution range)	99.0 (44.82-2.56) 99.0 (44.82-2.56)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.75 (at 2.58Å)	Xtriage
Refinement program	PHENIX 1.21.2_5419	Depositor
R, R_{free}	0.218 , 0.251 0.218 , 0.251	Depositor DCC
R_{free} test set	907 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	71.7	Xtriage
Anisotropy	0.219	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 56.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.034 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3402	wwPDB-VP
Average B, all atoms (Å ²)	78.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, A1JJH, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/1603	0.54	0/2191
2	B	0.41	0/1838	0.58	0/2505
All	All	0.41	0/3441	0.56	0/4696

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1561	0	1457	27	0
2	B	1794	0	1632	30	0
3	A	34	0	0	1	0
4	B	4	0	3	1	0
5	B	1	0	0	0	0
6	A	3	0	0	0	0
6	B	5	0	0	0	0
All	All	3402	0	3092	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:391:VAL:HB	1:A:530:LYS:HG2	1.69	0.73
2:B:311:ILE:HG23	2:B:313:LEU:H	1.52	0.72
2:B:168:MET:HE2	2:B:370:ASN:H	1.59	0.68
1:A:455:ILE:HD13	1:A:520:MET:HE1	1.80	0.63
2:B:198:TYR:HB3	2:B:324:ILE:HG22	1.81	0.61
2:B:234:TRP:CD2	2:B:326:LYS:HD2	2.37	0.60
2:B:192:GLU:HG3	2:B:234:TRP:CD1	2.42	0.54
2:B:255:ARG:HB2	4:B:401:ACT:H3	1.90	0.54
2:B:382:ASN:HA	2:B:385:LEU:HD11	1.90	0.53
2:B:342:ARG:HG2	2:B:378:PHE:CZ	2.44	0.53
2:B:168:MET:CE	2:B:370:ASN:H	2.23	0.51
1:A:410:THR:HG22	1:A:413:GLU:CD	2.35	0.51
2:B:228:ARG:HA	2:B:293:LYS:O	2.12	0.50
2:B:341:ARG:HD3	2:B:385:LEU:HB2	1.93	0.49
1:A:457:TRP:CE3	1:A:514:PRO:HD3	2.48	0.49
1:A:504:VAL:HG23	2:B:130:PRO:HB2	1.96	0.48
2:B:128:HIS:HB3	2:B:132:ASN:HD22	1.78	0.48
2:B:342:ARG:HG2	2:B:378:PHE:CE1	2.49	0.48
2:B:185:LYS:HE2	2:B:223:GLU:O	2.14	0.48
1:A:451:ARG:NH1	2:B:242:ASP:OD2	2.47	0.47
2:B:382:ASN:HA	2:B:385:LEU:CD1	2.44	0.47
2:B:332:HIS:CD2	2:B:336:HIS:NE2	2.83	0.47
1:A:370:PRO:HG2	1:A:557:LEU:HD23	1.96	0.46
1:A:368:LEU:C	1:A:370:PRO:HD2	2.41	0.46
1:A:373:TRP:HB2	1:A:551:LEU:HD23	1.98	0.46
2:B:157:ASP:HA	2:B:160:ILE:HD12	1.97	0.46
1:A:440:GLY:O	1:A:444:LEU:HB2	2.16	0.46
1:A:437:MET:HB2	2:B:255:ARG:NH1	2.31	0.45
1:A:383:VAL:HB	1:A:422:LEU:HD21	1.98	0.45
1:A:437:MET:HE2	1:A:437:MET:HB3	1.78	0.45
1:A:449:TYR:CD2	1:A:488:LYS:HB2	2.53	0.44
1:A:555:HIS:NE2	1:A:557:LEU:HD11	2.32	0.44
2:B:276:ASP:HB3	2:B:279:ALA:HB2	2.00	0.44
1:A:381:LEU:HD23	1:A:547:LEU:HD13	1.99	0.43
1:A:539:ASN:O	1:A:541:GLN:HG2	2.18	0.43
1:A:409:LEU:HD11	3:A:601:A1JJH:N26	2.34	0.43
1:A:411:ASP:O	1:A:415:ARG:HG3	2.19	0.42
2:B:171:GLN:HE21	2:B:171:GLN:HB3	1.68	0.42
2:B:332:HIS:HD2	2:B:336:HIS:NE2	2.18	0.42
2:B:337:PHE:HB3	2:B:339:LEU:HG	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:404:LEU:HD12	1:A:404:LEU:H	1.85	0.41
2:B:385:LEU:HD12	2:B:385:LEU:H	1.85	0.41
1:A:575:SER:O	1:A:576:LYS:C	2.63	0.41
2:B:261:TRP:CE2	2:B:327:PRO:HD3	2.56	0.41
1:A:568:ARG:O	1:A:568:ARG:HD3	2.21	0.41
2:B:245:ARG:HG3	2:B:289:LEU:HD21	2.02	0.41
2:B:192:GLU:HG3	2:B:234:TRP:HD1	1.85	0.41
2:B:164:ASN:OD1	2:B:356:GLY:HA3	2.21	0.40
1:A:393:MET:CG	1:A:429:PHE:HB2	2.51	0.40
1:A:436:ALA:HA	1:A:439:LEU:HB3	2.04	0.40
1:A:404:LEU:HD13	1:A:408:THR:HG21	2.03	0.40
1:A:478:HIS:HE1	2:B:257:GLU:OE2	2.04	0.40
1:A:393:MET:HG3	1:A:429:PHE:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	199/228 (87%)	187 (94%)	10 (5%)	2 (1%)	12	17
2	B	228/307 (74%)	222 (97%)	6 (3%)	0	100	100
All	All	427/535 (80%)	409 (96%)	16 (4%)	2 (0%)	24	33

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	401	HIS
1	A	573	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	156/200 (78%)	153 (98%)	3 (2%)	50	65
2	B	173/274 (63%)	165 (95%)	8 (5%)	24	35
All	All	329/474 (69%)	318 (97%)	11 (3%)	33	48

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	444	LEU
1	A	464	GLN
1	A	498	LEU
2	B	179	ILE
2	B	183	THR
2	B	185	LYS
2	B	191	LEU
2	B	255	ARG
2	B	256	CYS
2	B	332	HIS
2	B	364	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	478	HIS
1	A	541	GLN
2	B	128	HIS
2	B	171	GLN
2	B	332	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ACT	B	401	-	3,3,3	1.23	0	3,3,3	1.06	0
3	A1JJH	A	601	-	37,38,38	3.19	20 (54%)	47,57,57	3.11	23 (48%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A1JJH	A	601	-	-	3/12/50/50	0/5/5/5

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601	A1JJH	C18-C17	7.09	1.61	1.55
3	A	601	A1JJH	C14-C12	6.60	1.48	1.39
3	A	601	A1JJH	C30-C31	5.98	1.59	1.51
3	A	601	A1JJH	C14-N15	5.00	1.52	1.41
3	A	601	A1JJH	C18-N20	4.85	1.54	1.48
3	A	601	A1JJH	C16-C17	4.83	1.58	1.53
3	A	601	A1JJH	C21-N20	4.57	1.47	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601	A1JJH	C19-N20	4.34	1.54	1.48
3	A	601	A1JJH	C19-C17	3.63	1.58	1.55
3	A	601	A1JJH	C25-N26	3.55	1.43	1.36
3	A	601	A1JJH	C11-C12	3.51	1.43	1.38
3	A	601	A1JJH	C27-N26	3.50	1.50	1.45
3	A	601	A1JJH	C25-N24	3.23	1.40	1.34
3	A	601	A1JJH	C07-C08	3.20	1.57	1.51
3	A	601	A1JJH	C21-N22	3.18	1.41	1.34
3	A	601	A1JJH	C28-C21	3.06	1.44	1.39
3	A	601	A1JJH	C30-N15	2.90	1.51	1.46
3	A	601	A1JJH	C16-N15	2.80	1.48	1.46
3	A	601	A1JJH	C28-C25	2.56	1.46	1.39
3	A	601	A1JJH	C31-N29	2.42	1.37	1.34

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	A1JJH	C25-C28-C21	9.55	124.28	116.64
3	A	601	A1JJH	C10-C14-C12	-7.39	112.03	119.94
3	A	601	A1JJH	C30-N15-C16	-6.40	103.79	112.68
3	A	601	A1JJH	C27-N26-C25	5.66	128.20	122.96
3	A	601	A1JJH	C28-C21-N22	-4.97	114.56	122.82
3	A	601	A1JJH	C17-C19-N20	4.57	93.28	87.90
3	A	601	A1JJH	C28-C25-N24	-4.49	117.03	122.92
3	A	601	A1JJH	C04-C05-N06	3.68	114.92	111.16
3	A	601	A1JJH	C10-C14-N15	3.53	127.95	122.42
3	A	601	A1JJH	C09-C10-C14	3.51	126.13	119.10
3	A	601	A1JJH	C17-C18-N20	3.34	91.84	87.90
3	A	601	A1JJH	C12-C11-C08	3.21	124.14	120.81
3	A	601	A1JJH	C23-N22-C21	3.09	117.55	114.95
3	A	601	A1JJH	C19-N20-C21	3.08	135.20	125.17
3	A	601	A1JJH	C03-C02-C34	-3.05	105.65	110.04
3	A	601	A1JJH	C09-C08-C11	-3.02	114.38	118.55
3	A	601	A1JJH	C19-N20-C18	-2.81	89.11	93.99
3	A	601	A1JJH	C28-C21-N20	2.80	125.73	122.35
3	A	601	A1JJH	C34-C33-N06	2.66	113.87	111.16
3	A	601	A1JJH	C07-C08-C09	2.50	125.36	120.75
3	A	601	A1JJH	C18-N20-C21	2.47	133.20	125.17
3	A	601	A1JJH	O32-C31-N29	-2.13	118.83	121.49
3	A	601	A1JJH	N22-C21-N20	2.06	119.71	116.76

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	601	A1JJH	C28-C25-N26-C27
3	A	601	A1JJH	N24-C25-N26-C27
3	A	601	A1JJH	C12-C14-N15-C16

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	401	ACT	1	0
3	A	601	A1JJH	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	203/228 (89%)	0.64	12 (5%) 28 28	56, 80, 113, 124	0
2	B	238/307 (77%)	0.57	18 (7%) 20 19	50, 72, 110, 125	0
All	All	441/535 (82%)	0.60	30 (6%) 23 23	50, 75, 111, 125	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	509	SER	4.0
2	B	125	ASP	3.8
1	A	507	VAL	3.2
2	B	120	CYS	3.2
1	A	508	ARG	3.1
1	A	386	LEU	3.0
2	B	152	LEU	2.9
2	B	134	ILE	2.9
2	B	122	HIS	2.8
2	B	381	PRO	2.7
2	B	310	ASP	2.6
2	B	137	VAL	2.6
1	A	575	SER	2.6
2	B	179	ILE	2.5
2	B	126	THR	2.4
1	A	576	LYS	2.4
1	A	475	TRP	2.4
2	B	258	ASP	2.3
2	B	390	GLU	2.3
1	A	512	HIS	2.2
2	B	119	TYR	2.2
2	B	339	LEU	2.2
1	A	574	ILE	2.2
1	A	550	GLN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	485	VAL	2.1
2	B	202	THR	2.1
2	B	313	LEU	2.1
1	A	368	LEU	2.1
2	B	389	THR	2.1
2	B	377	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	A1JJH	A	601	34/34	0.88	0.12	75,83,91,94	0
4	ACT	B	401	4/4	0.94	0.18	68,71,75,81	0
5	MG	B	402	1/1	0.96	0.14	62,62,62,62	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.