



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 9, 2026 – 09:46 PM UTC

PDB ID : 9T3L / pdb_00009t3l
Title : Crystal structure of the Acl1 ankyrin repeat domain in complex with the second Rpl1 domain
Authors : Burchert, F.; Kiontke, S.; Bange, G.
Deposited on : 2025-10-28
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1091 (2.54-2.54)
Clashscore	190562	1120 (2.54-2.54)
Ramachandran outliers	187476	1106 (2.54-2.54)
Sidechain outliers	187428	1106 (2.54-2.54)
RSRZ outliers	180081	1091 (2.54-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	104	<div> <div>4%</div> <div>63%</div> <div>21%</div> <div>15%</div> </div>
2	B	128	<div> <div>%</div> <div>84%</div> <div>14%</div> <div>.</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1663 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Large ribosomal subunit protein uL1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	88	Total	C	N	O	S	0	0	0
			679	435	116	125	3			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	159	GLY	-	expression tag	UNP P0CX43
A	160	SER	-	expression tag	UNP P0CX43
A	161	HIS	-	expression tag	UNP P0CX43
A	162	HIS	-	expression tag	UNP P0CX43
A	163	HIS	-	expression tag	UNP P0CX43
A	164	HIS	-	expression tag	UNP P0CX43
A	165	HIS	-	expression tag	UNP P0CX43
A	166	HIS	-	expression tag	UNP P0CX43

- Molecule 2 is a protein called Ankyrin repeat-containing protein YCR051W.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	126	Total	C	N	O	S	0	0	0
			975	601	169	200	5			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	128	SER	-	expression tag	UNP P25631

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	O	0	0
			1	1		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	8	Total	O	0	0
			8	8		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Large ribosomal subunit protein uL1A



- Molecule 2: Ankyrin repeat-containing protein YCR051W



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	101.65Å 101.65Å 108.13Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.07 – 2.55 46.07 – 2.55	Depositor EDS
% Data completeness (in resolution range)	100.0 (46.07-2.55) 100.0 (46.07-2.55)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.18 (at 2.54Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.195 , 0.242 0.194 , 0.242	Depositor DCC
R_{free} test set	563 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	47.5	Xtriage
Anisotropy	0.048	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 49.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	1663	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/689	0.66	0/924
2	B	0.49	0/992	0.65	0/1344
All	All	0.48	0/1681	0.65	0/2268

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	679	0	717	16	0
2	B	975	0	919	11	0
3	A	1	0	0	0	0
3	B	8	0	0	0	0
All	All	1663	0	1636	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:81:VAL:HG21	2:B:117:GLU:HG3	1.45	0.96

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:136:THR:HG21	1:A:151:VAL:HG11	1.63	0.80
1:A:73:ASP:OD1	1:A:76:ARG:NH1	2.16	0.78
1:A:95:LYS:HE2	1:A:122:ARG:O	1.90	0.72
1:A:138:VAL:HG23	1:A:147:LYS:HG3	1.74	0.69
1:A:67:ILE:HD13	1:A:111:ILE:HB	1.84	0.60
2:B:86:LEU:HD13	2:B:91:LEU:HD23	1.86	0.58
2:B:115:ASP:O	2:B:119:ILE:HD12	2.03	0.58
1:A:90:LEU:HD22	1:A:124:LEU:HD21	1.86	0.57
1:A:65:ILE:HG13	1:A:148:VAL:HG13	1.87	0.56
1:A:67:ILE:HD12	1:A:144:LEU:HD11	1.91	0.52
1:A:143:ASP:OD2	1:A:146:GLY:N	2.36	0.49
2:B:115:ASP:OD2	2:B:117:GLU:HB3	2.12	0.49
1:A:111:ILE:CD1	1:A:148:VAL:HG22	2.43	0.48
1:A:76:ARG:NH2	1:A:142:ASP:O	2.46	0.48
1:A:72:PHE:CE2	1:A:76:ARG:HD2	2.49	0.47
1:A:104:SER:O	1:A:133:LYS:NZ	2.38	0.47
1:A:144:LEU:O	1:A:148:VAL:HG23	2.15	0.47
2:B:84:ALA:O	2:B:88:VAL:HG23	2.16	0.46
2:B:52:LEU:HG	2:B:56:MET:HE2	1.97	0.45
2:B:125:LYS:HD3	2:B:125:LYS:HA	1.65	0.45
1:A:119:GLN:HG2	1:A:122:ARG:NH1	2.32	0.45
2:B:95:PHE:HB2	2:B:122:MET:HE1	2.00	0.44
2:B:76:HIS:CE1	2:B:105:PRO:HD3	2.55	0.42
1:A:148:VAL:HA	1:A:151:VAL:HG22	2.02	0.41
2:B:44:ALA:HB3	2:B:53:LEU:HD13	2.02	0.41
2:B:105:PRO:HB2	2:B:122:MET:HE2	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	86/104 (83%)	83 (96%)	3 (4%)	0	100	100
2	B	124/128 (97%)	124 (100%)	0	0	100	100
All	All	210/232 (90%)	207 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	77/92 (84%)	77 (100%)	0	100	100
2	B	104/106 (98%)	103 (99%)	1 (1%)	68	80
All	All	181/198 (91%)	180 (99%)	1 (1%)	78	87

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	15	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	88/104 (84%)	0.21	4 (4%) 38 38	40, 65, 109, 160	0
2	B	126/128 (98%)	-0.25	1 (0%) 82 84	36, 56, 88, 101	0
All	All	214/232 (92%)	-0.06	5 (2%) 61 62	36, 61, 96, 160	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	145	TYR	6.8
2	B	127	GLY	4.4
1	A	148	VAL	3.6
1	A	149	THR	2.8
1	A	144	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.