



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 4, 2026 – 09:42 PM UTC

PDB ID : 9U64 / pdb_00009u64
Title : Crystal structure of human glutaminyl cyclase in complex with Inhibitor CL21
Authors : Li, G.-B.; Meng, F.-B.; Chen, Y.-T.
Deposited on : 2025-03-22
Resolution : 2.27 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1898 (2.26-2.26)
Clashscore	190562	2005 (2.26-2.26)
Ramachandran outliers	187476	1965 (2.26-2.26)
Sidechain outliers	187428	1966 (2.26-2.26)
RSRZ outliers	180081	1898 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	361	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	A	407	-	X	X	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	A	408	-	-	X	-
5	GOL	A	409	-	-	-	X

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 3035 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutaminyl-peptide cyclotransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	329	Total	C	N	O	S	0	3	0
			2684	1715	467	493	9			


- Molecule 2 is 3-(2,4-dimethoxyphenyl)-6-[(5-methylimidazol-1-yl)methyl]-1H-pyridin-2-one (CCD ID: A1EN4) (formula: C₁₈H₁₉N₃O₃) (labeled as "Ligand of Interest" by depositor).



- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	3	Total	Mg	0	0
			3	3		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	299	Total	O	0	0
			299	299		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutaminyl-peptide cyclotransferase



4 Data and refinement statistics

Property	Value	Source
Space group	F 41 3 2	Depositor
Cell constants a, b, c, α , β , γ	277.71 Å 277.71 Å 277.71 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.91 – 2.27 43.91 – 2.27	Depositor EDS
% Data completeness (in resolution range)	99.6 (43.91-2.27) 99.5 (43.91-2.27)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.17 (at 2.27 Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.178 , 0.210 0.191 , 0.224	Depositor DCC
R_{free} test set	2000 reflections (4.65%)	wwPDB-VP
Wilson B-factor (Å ²)	32.2	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 52.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3035	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, A1EN4, ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.22	4/2762 (0.1%)	1.11	1/3758 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	107	ASP	CG-OD2	6.37	1.37	1.25
1	A	302	VAL	CB-CG1	6.17	1.73	1.52
1	A	124	ILE	CG1-CD1	-6.01	1.28	1.51
1	A	188	LYS	CA-C	-5.55	1.45	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	107	ASP	CA-CB-CG	5.12	117.72	112.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	187	SER	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2684	0	2603	30	0
2	A	24	0	0	0	0
3	A	1	0	0	0	0
4	A	3	0	0	0	0
5	A	24	0	31	10	0
6	A	299	0	0	22	1
All	All	3035	0	2634	35	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:97[A]:ARG:NH2	6:A:502:HOH:O	1.84	1.10
1:A:49:ASN:OD1	6:A:501:HOH:O	1.77	1.00
1:A:54:ARG:NH1	6:A:505:HOH:O	1.99	0.95
1:A:301:GLY:O	6:A:503:HOH:O	1.89	0.90
1:A:107:ASP:OD1	6:A:504:HOH:O	1.97	0.80
1:A:312:ARG:HH21	5:A:407:GOL:H11	1.50	0.76
1:A:309:PRO:HB3	5:A:407:GOL:H12	1.70	0.72
1:A:233:ARG:NH1	6:A:507:HOH:O	2.23	0.70
5:A:408:GOL:H11	6:A:750:HOH:O	1.91	0.68
1:A:302:VAL:HG13	6:A:652:HOH:O	1.96	0.65
1:A:263:ASN:HB2	6:A:742:HOH:O	2.01	0.59
1:A:93[B]:GLN:NE2	6:A:512:HOH:O	2.36	0.58
1:A:128:ASN:ND2	6:A:513:HOH:O	2.37	0.58
1:A:263:ASN:C	6:A:510:HOH:O	2.48	0.56
1:A:49:ASN:HB2	6:A:712:HOH:O	2.03	0.56
1:A:302:VAL:HA	6:A:652:HOH:O	2.08	0.54
1:A:131:ALA:HB3	1:A:134:HIS:CE1	2.47	0.50
1:A:298:SER:OG	6:A:506:HOH:O	2.19	0.50
1:A:101:ASP:HB2	1:A:182:LYS:HE3	1.93	0.50
5:A:408:GOL:H11	6:A:546:HOH:O	2.12	0.49
1:A:188:LYS:O	1:A:188:LYS:CG	2.58	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:309:PRO:CB	5:A:407:GOL:H12	2.42	0.48
1:A:188:LYS:O	1:A:189:PRO:C	2.55	0.47
1:A:312:ARG:HD3	5:A:407:GOL:H11	1.97	0.47
1:A:182:LYS:HD3	6:A:568:HOH:O	2.15	0.46
5:A:408:GOL:C1	6:A:546:HOH:O	2.62	0.46
1:A:258:PRO:HB3	1:A:297:TYR:CE1	2.51	0.46
1:A:40:LYS:NZ	6:A:524:HOH:O	2.49	0.45
5:A:408:GOL:C3	6:A:750:HOH:O	2.64	0.45
1:A:33:ALA:HB2	6:A:528:HOH:O	2.17	0.44
5:A:408:GOL:H31	6:A:750:HOH:O	2.17	0.43
1:A:308:ILE:O	1:A:312:ARG:HG2	2.20	0.42
1:A:186:ASP:C	1:A:188:LYS:HG2	2.45	0.41
1:A:308:ILE:HB	1:A:309:PRO:HD3	2.01	0.41
1:A:54:ARG:HE	5:A:406:GOL:H32	1.86	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:718:HOH:O	6:A:760:HOH:O[59_555]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	330/361 (91%)	315 (96%)	12 (4%)	3 (1%)	14 12

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	212	SER
1	A	300	GLY

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	188	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	293/314 (93%)	283 (97%)	10 (3%)	32	40

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	SER
1	A	151	ASN
1	A	176	LYS
1	A	184	VAL
1	A	188	LYS
1	A	210	GLN
1	A	242	ASP
1	A	284	LEU
1	A	288	SER
1	A	295	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	44	GLN
1	A	55	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	A1EN4	A	401	3	25,26,26	3.17	11 (44%)	26,36,36	2.81	11 (42%)
5	GOL	A	409	-	5,5,5	0.15	0	5,5,5	0.30	0
5	GOL	A	406	-	5,5,5	0.51	0	5,5,5	2.18	3 (60%)
5	GOL	A	407	-	5,5,5	1.18	0	5,5,5	3.68	4 (80%)
5	GOL	A	408	-	5,5,5	1.19	0	5,5,5	1.48	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1EN4	A	401	3	-	7/12/12/12	0/3/3/3
5	GOL	A	409	-	-	4/4/4/4	-
5	GOL	A	406	-	-	0/4/4/4	-
5	GOL	A	407	-	-	3/4/4/4	-
5	GOL	A	408	-	-	2/4/4/4	-

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	A1EN4	C16-N24	7.01	1.47	1.37
2	A	401	A1EN4	C15-C16	6.49	1.52	1.37
2	A	401	A1EN4	C14-C03	6.05	1.48	1.37
2	A	401	A1EN4	C22-N18	-5.09	1.30	1.38
2	A	401	A1EN4	C02-N24	4.37	1.47	1.38
2	A	401	A1EN4	O01-C02	-3.94	1.16	1.23
2	A	401	A1EN4	C03-C02	2.87	1.49	1.46
2	A	401	A1EN4	C15-C14	2.79	1.48	1.41
2	A	401	A1EN4	C13-C04	-2.75	1.35	1.39
2	A	401	A1EN4	C21-N20	-2.73	1.32	1.37
2	A	401	A1EN4	C19-N18	-2.22	1.32	1.36

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	A1EN4	O01-C02-C03	-7.85	115.21	125.19
5	A	407	GOL	C3-C2-C1	5.22	130.95	111.80
2	A	401	A1EN4	C04-C03-C02	-4.92	114.10	120.02
2	A	401	A1EN4	C15-C14-C03	-4.69	118.35	123.35
2	A	401	A1EN4	C13-C12-C09	4.31	124.66	119.73
5	A	407	GOL	O3-C3-C2	4.15	129.08	110.38
2	A	401	A1EN4	C13-C04-C03	-3.88	113.55	119.60
5	A	407	GOL	O1-C1-C2	3.83	127.60	110.38
2	A	401	A1EN4	C05-C04-C03	3.54	126.23	121.56
5	A	406	GOL	O3-C3-C2	-2.89	97.37	110.38
5	A	407	GOL	O2-C2-C1	2.74	120.50	109.18
5	A	408	GOL	O1-C1-C2	2.66	122.36	110.38
5	A	406	GOL	O2-C2-C1	2.63	120.09	109.18
5	A	406	GOL	C3-C2-C1	-2.46	102.76	111.80
2	A	401	A1EN4	C23-C22-N18	2.46	125.72	122.80
2	A	401	A1EN4	C14-C03-C02	2.43	120.77	119.39
2	A	401	A1EN4	O06-C05-C08	-2.33	120.07	124.08
2	A	401	A1EN4	C04-C03-C14	2.12	125.03	122.32
2	A	401	A1EN4	O06-C05-C04	2.01	119.48	116.55

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	A1EN4	N24-C16-C17-N18
5	A	407	GOL	O1-C1-C2-C3
5	A	408	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
5	A	409	GOL	O1-C1-C2-C3
5	A	409	GOL	C1-C2-C3-O3
2	A	401	A1EN4	C08-C09-O10-C11
2	A	401	A1EN4	C12-C09-O10-C11
2	A	401	A1EN4	C08-C05-O06-C07
5	A	407	GOL	O2-C2-C3-O3
5	A	408	GOL	O1-C1-C2-O2
5	A	409	GOL	O2-C2-C3-O3
2	A	401	A1EN4	C04-C05-O06-C07
5	A	409	GOL	O1-C1-C2-O2
5	A	407	GOL	O1-C1-C2-O2
2	A	401	A1EN4	C15-C16-C17-N18
2	A	401	A1EN4	C14-C03-C04-C13

There are no ring outliers.

3 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	406	GOL	1	0
5	A	407	GOL	4	0
5	A	408	GOL	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	329/361 (91%)	0.00	24 (7%) 21 19	17, 29, 64, 115	3 (0%)

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	33	ALA	10.9
1	A	34	SER	7.8
1	A	184	VAL	6.9
1	A	188	LYS	5.1
1	A	187	SER	5.1
1	A	131	ALA	4.5
1	A	149	TRP	4.2
1	A	112	GLN	4.1
1	A	183	THR	4.0
1	A	117	TYR	3.5
1	A	35	ALA	3.5
1	A	297	TYR	3.4
1	A	185	SER	3.3
1	A	150	ASN	3.1
1	A	233	ARG	3.1
1	A	115	TYR	3.0
1	A	302	VAL	3.0
1	A	182	LYS	2.9
1	A	97[A]	ARG	2.8
1	A	186	ASP	2.6
1	A	301	GLY	2.5
1	A	269	GLU	2.1
1	A	300	GLY	2.1
1	A	93[A]	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	A	409	6/6	0.57	0.46	12,30,41,64	0
5	GOL	A	406	6/6	0.76	0.25	48,67,70,73	0
5	GOL	A	408	6/6	0.80	0.23	33,58,76,78	0
4	MG	A	404	1/1	0.85	0.39	67,67,67,67	0
2	A1EN4	A	401	24/24	0.86	0.13	17,25,36,38	0
5	GOL	A	407	6/6	0.90	0.24	30,43,64,72	0
4	MG	A	405	1/1	0.96	0.12	67,67,67,67	0
4	MG	A	403	1/1	0.99	0.05	20,20,20,20	0
3	ZN	A	402	1/1	1.00	0.02	22,22,22,22	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1EN4 A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.