



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 9, 2026 – 10:29 PM UTC

PDB ID : 9UAI / pdb\_00009uai  
Title : Crystal structure of cyanobacterial CRD of CmpBCD in the presence of nitrate  
Authors : Li, Q.Y.; Li, B.; Zhou, C.Z.; Jiang, Y.L.  
Deposited on : 2025-04-01  
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	400	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3187 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Nitrate transport ATP-binding subunits C and D.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	373	Total	C	N	O	S	0	0	0
			2915	1851	516	531	17			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	260	MET	-	initiating methionine	UNP A0A2Z5WYP5
E	261	GLY	-	expression tag	UNP A0A2Z5WYP5
E	262	SER	-	expression tag	UNP A0A2Z5WYP5
E	263	SER	-	expression tag	UNP A0A2Z5WYP5
E	264	HIS	-	expression tag	UNP A0A2Z5WYP5
E	265	HIS	-	expression tag	UNP A0A2Z5WYP5
E	266	HIS	-	expression tag	UNP A0A2Z5WYP5
E	267	HIS	-	expression tag	UNP A0A2Z5WYP5
E	268	HIS	-	expression tag	UNP A0A2Z5WYP5
E	269	HIS	-	expression tag	UNP A0A2Z5WYP5
E	270	SER	-	expression tag	UNP A0A2Z5WYP5
E	271	SER	-	expression tag	UNP A0A2Z5WYP5
E	272	GLY	-	expression tag	UNP A0A2Z5WYP5
E	273	LYS	-	expression tag	UNP A0A2Z5WYP5
E	349	ALA	ASN	conflict	UNP A0A2Z5WYP5
E	391	SER	ASN	conflict	UNP A0A2Z5WYP5
E	396	GLN	THR	conflict	UNP A0A2Z5WYP5
E	425	LEU	HIS	conflict	UNP A0A2Z5WYP5

- Molecule 2 is NITRATE ION (CCD ID: NO3) (formula: NO<sub>3</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	E	1	Total	N	O	0	0
			4	1	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	E	268	Total	O	0	0
			268	268		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitrate transport ATP-binding subunits C and D



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	92.27Å 49.96Å 97.58Å 90.00° 116.70° 90.00°	Depositor
Resolution (Å)	13.07 – 1.75 13.07 – 1.75	Depositor EDS
% Data completeness (in resolution range)	99.5 (13.07-1.75) 99.3 (13.07-1.75)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.56 (at 1.75Å)	Xtriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, $R_{free}$	0.197 , 0.229 0.197 , 0.230	Depositor DCC
$R_{free}$ test set	1999 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.0	Xtriage
Anisotropy	0.814	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.44 , 52.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.015 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3187	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.79% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	E	0.35	0/2989	0.60	1/4066 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	621	SER	CB-CA-C	-5.63	109.57	117.23

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	2915	0	2855	38	0
2	E	4	0	0	1	0
3	E	268	0	0	12	1
All	All	3187	0	2855	38	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:281:LEU:HD13	1:E:311:VAL:HG22	1.60	0.84
1:E:386:LYS:NZ	1:E:390:GLN:OE1	2.13	0.80
1:E:598:GLU:OE1	3:E:801:HOH:O	2.03	0.74
1:E:352:CYS:HB3	3:E:928:HOH:O	1.90	0.70
1:E:393:ARG:HH12	1:E:421:ILE:HA	1.60	0.66
1:E:320:ARG:NH1	3:E:804:HOH:O	2.30	0.62
1:E:422:ASP:OD1	1:E:649:ARG:NH2	2.42	0.52
1:E:275:GLY:O	3:E:803:HOH:O	2.19	0.51
1:E:422:ASP:CG	1:E:649:ARG:HH22	2.19	0.51
1:E:518:GLU:OE2	1:E:521:ARG:NH1	2.43	0.51
1:E:555:GLU:HB3	1:E:556:PRO:HD2	1.93	0.50
1:E:541:ASN:ND2	1:E:555:GLU:OE1	2.44	0.50
1:E:407:MET:HA	1:E:407:MET:HE2	1.93	0.49
1:E:539:LEU:HA	3:E:957:HOH:O	2.12	0.49
1:E:281:LEU:HD11	1:E:308:LEU:HD22	1.95	0.49
1:E:626:GLN:NE2	1:E:630:ASP:OD1	2.47	0.48
1:E:622:ARG:CG	3:E:874:HOH:O	2.62	0.48
1:E:351:ALA:O	1:E:353:LYS:HE3	2.15	0.47
1:E:402:VAL:HG11	2:E:701:NO3:O3	2.15	0.47
1:E:300:LYS:HD2	3:E:861:HOH:O	2.15	0.47
1:E:278:LYS:HD3	1:E:280:ASN:O	2.15	0.46
1:E:402:VAL:HG22	3:E:827:HOH:O	2.17	0.45
1:E:509:MET:HG2	1:E:562:TYR:CE1	2.52	0.44
1:E:619:THR:N	3:E:802:HOH:O	2.46	0.43
1:E:509:MET:SD	1:E:615:LEU:HD11	2.59	0.43
1:E:457:PHE:CE1	1:E:556:PRO:HD3	2.54	0.42
1:E:458:ARG:HG3	1:E:536:TYR:CE2	2.54	0.42
1:E:453:GLU:HB3	1:E:454:PRO:HA	2.02	0.42
1:E:558:HIS:ND1	1:E:558:HIS:C	2.77	0.42
1:E:377:ARG:HE	1:E:377:ARG:HB3	1.43	0.42
1:E:622:ARG:HG3	3:E:874:HOH:O	2.18	0.42
1:E:353:LYS:HE2	1:E:353:LYS:HB3	1.76	0.41
1:E:344:LEU:HD22	1:E:353:LYS:HD2	2.03	0.41
1:E:509:MET:HG2	1:E:562:TYR:CD1	2.56	0.41
1:E:300:LYS:HE3	1:E:519:GLU:OE2	2.21	0.41
1:E:622:ARG:HG2	3:E:874:HOH:O	2.20	0.41
1:E:342:MET:HG3	1:E:578:ILE:CG1	2.51	0.41
1:E:620:TYR:N	3:E:802:HOH:O	2.12	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:992:HOH:O	3:E:1042:HOH:O[3_545]	1.97	0.23

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	369/400 (92%)	356 (96%)	13 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	306/330 (93%)	304 (99%)	2 (1%)	76	67

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	338	SER
1	E	630	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	436	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	E	516	ASN
1	E	644	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NO3	E	701	-	1,3,3	0.51	0	0,3,3	-	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	701	NO3	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	E	373/400 (93%)	0.19	30 (8%) 18 22	10, 21, 43, 96	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	623	ASN	6.5
1	E	616	THR	6.3
1	E	624	ALA	6.3
1	E	542	PRO	5.1
1	E	274	HIS	4.7
1	E	621	SER	4.4
1	E	556	PRO	4.0
1	E	540	GLY	3.9
1	E	618	ILE	3.8
1	E	541	ASN	3.7
1	E	558	HIS	3.6
1	E	644	HIS	3.6
1	E	657	ALA	3.4
1	E	622	ARG	3.1
1	E	392	THR	3.1
1	E	393	ARG	3.1
1	E	619	THR	3.0
1	E	275	GLY	3.0
1	E	557	ALA	2.8
1	E	353	LYS	2.8
1	E	543	ASP	2.6
1	E	352	CYS	2.5
1	E	351	ALA	2.5
1	E	394	ASP	2.3
1	E	617	ASP	2.3
1	E	330	TYR	2.3
1	E	630	ASP	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	E	555	GLU	2.1
1	E	620	TYR	2.1
1	E	631	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NO3	E	701	4/4	0.89	0.13	21,26,29,30	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around NO3 E 701:**

$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.