



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 21, 2026 – 03:07 AM JST

PDB ID : 9ULQ / pdb_00009ulq
Title : Recombinant Human Serum Albumin protein 3
Authors : Xiang, W.; Yue, Z.L.; Su, J.Y.
Deposited on : 2025-04-21
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1190 (2.46-2.46)
Clashscore	190562	1229 (2.46-2.46)
Ramachandran outliers	187476	1218 (2.46-2.46)
Sidechain outliers	187428	1218 (2.46-2.46)
RSRZ outliers	180081	1190 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	585	<div> <div>13%</div> <div>84%</div> <div>13%</div> <div>.</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4626 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Albumin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	570	Total	C	N	O	S	0	0	0
			4538	2866	768	864	40			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	88	Total	O	0	0
			88	88		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Albumin



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	56.09Å 87.44Å 58.87Å 90.00° 104.76° 90.00°	Depositor
Resolution (Å)	19.94 – 2.45 19.94 – 2.45	Depositor EDS
% Data completeness (in resolution range)	96.9 (19.94-2.45) 96.7 (19.94-2.45)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.16 (at 2.44Å)	Xtriage
Refinement program	PHENIX (1.21.1_5286: ???)	Depositor
R, R_{free}	0.246 , 0.287 0.247 , 0.287	Depositor DCC
R_{free} test set	1951 reflections (9.62%)	wwPDB-VP
Wilson B-factor (Å ²)	31.3	Xtriage
Anisotropy	0.211	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 36.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.036 for l,-k,h	Xtriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	4626	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.15	0/4626	0.35	0/6238

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4538	0	4461	47	0
2	A	88	0	0	1	0
All	All	4626	0	4461	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:132:GLU:HG3	1:A:136:LYS:HE2	1.73	0.70
1:A:30:TYR:HD2	1:A:102:PHE:HB3	1.61	0.65
1:A:153:GLU:OE2	1:A:288:HIS:ND1	2.31	0.63
1:A:314:ASP:HB3	1:A:317:LYS:HE2	1.81	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:224:PRO:HD2	1:A:296:ASP:HB3	1.83	0.60
1:A:305:LEU:HD12	1:A:309:PHE:HD2	1.67	0.60
1:A:305:LEU:HD12	1:A:309:PHE:CD2	2.37	0.59
1:A:57:GLU:H	1:A:57:GLU:CD	2.11	0.58
1:A:433:VAL:HG22	1:A:452:TYR:CD1	2.39	0.58
1:A:31:LEU:HD12	1:A:74:LEU:HG	1.86	0.56
1:A:394:LEU:O	1:A:398:LEU:HB2	2.06	0.56
1:A:30:TYR:CD2	1:A:102:PHE:HB3	2.40	0.55
1:A:118:PRO:HD2	1:A:123:MET:HE3	1.88	0.55
1:A:208:GLU:OE2	1:A:212:LYS:NZ	2.39	0.54
1:A:433:VAL:HG22	1:A:452:TYR:HD1	1.72	0.54
1:A:365:ASP:HB2	1:A:368:GLU:OE2	2.06	0.53
1:A:564:LYS:O	1:A:566:THR:N	2.40	0.53
1:A:214:TRP:CD1	1:A:343:VAL:HG11	2.45	0.52
1:A:525:LYS:HE2	1:A:548:MET:HG3	1.92	0.52
1:A:9:HIS:HA	1:A:12:LYS:HE2	1.92	0.52
1:A:135:LEU:HD21	1:A:162:LYS:HB2	1.91	0.51
1:A:305:LEU:HD11	1:A:333:GLU:HB3	1.91	0.51
1:A:518:GLU:CD	1:A:521:ARG:HE	2.19	0.50
1:A:529:LEU:HD13	1:A:548:MET:HE3	1.92	0.50
1:A:532:LEU:HD11	1:A:548:MET:HE1	1.96	0.48
1:A:485:ARG:HB3	1:A:486:PRO:HD3	1.95	0.48
1:A:532:LEU:HD21	1:A:548:MET:HE1	1.95	0.47
1:A:198:LEU:O	1:A:202:SER:OG	2.32	0.47
1:A:503:ASN:OD1	1:A:505:GLU:N	2.49	0.46
1:A:174:LYS:H	1:A:174:LYS:HG3	1.63	0.45
1:A:285:GLU:OE1	1:A:285:GLU:N	2.44	0.44
1:A:156:PHE:CZ	1:A:160:ARG:HD2	2.52	0.44
1:A:408:LEU:HD13	1:A:427:SER:HB2	2.00	0.44
1:A:383:GLU:HB3	1:A:384:PRO:HD3	1.99	0.43
1:A:347:LEU:O	1:A:351:LYS:HG2	2.19	0.42
1:A:410:ARG:HG2	1:A:414:LYS:HE2	2.00	0.42
1:A:503:ASN:OD1	1:A:503:ASN:C	2.62	0.42
1:A:257:ARG:CZ	1:A:287:SER:HB3	2.49	0.42
1:A:29:GLN:HG2	1:A:147:PRO:HA	2.01	0.41
1:A:277:GLU:O	1:A:281:LYS:HG2	2.19	0.41
1:A:398:LEU:HD23	1:A:403:PHE:HA	2.02	0.41
1:A:384:PRO:O	1:A:388:ILE:HG12	2.21	0.41
1:A:95:GLU:OE2	1:A:98:ARG:NH2	2.50	0.41
1:A:429:ASN:ND2	2:A:614:HOH:O	2.46	0.41
1:A:378:LYS:HB3	1:A:379:PRO:HD3	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:564:LYS:HD3	1:A:565:GLU:HG2	2.03	0.41
1:A:66:LEU:HB3	1:A:70:PHE:CE2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	566/585 (97%)	543 (96%)	19 (3%)	4 (1%)	18 24

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	111	ASN
1	A	300	ALA
1	A	565	GLU
1	A	301	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	500/511 (98%)	496 (99%)	4 (1%)	73 83

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	245	CYS
1	A	324	ASP
1	A	347	LEU
1	A	515	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	104	GLN
1	A	318	ASN
1	A	397	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	570/585 (97%)	1.07	77 (13%) 7 5	15, 37, 77, 105	0

All (77) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	442	GLU	6.4
1	A	578	ALA	5.9
1	A	582	ALA	5.9
1	A	553	ALA	5.8
1	A	581	ALA	5.2
1	A	509	PHE	5.2
1	A	3	HIS	4.4
1	A	502	PHE	4.3
1	A	577	ALA	4.2
1	A	441	PRO	4.0
1	A	507	PHE	3.8
1	A	365	ASP	3.7
1	A	520	GLU	3.6
1	A	443	ALA	3.5
1	A	164	ALA	3.5
1	A	576	VAL	3.3
1	A	551	PHE	3.3
1	A	170	GLN	3.2
1	A	173	ASP	3.2
1	A	508	THR	3.2
1	A	535	HIS	3.1
1	A	302	LEU	3.0
1	A	571	GLU	2.9
1	A	300	ALA	2.9
1	A	388	ILE	2.8
1	A	347	LEU	2.8
1	A	367	HIS	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	552	ALA	2.8
1	A	330	PHE	2.8
1	A	517	SER	2.8
1	A	89	ASP	2.7
1	A	565	GLU	2.7
1	A	134	PHE	2.7
1	A	568	PHE	2.6
1	A	510	HIS	2.6
1	A	311	GLU	2.6
1	A	544	LEU	2.6
1	A	504	ALA	2.6
1	A	247	HIS	2.5
1	A	356	THR	2.5
1	A	493	VAL	2.5
1	A	579	SER	2.5
1	A	369	CYS	2.4
1	A	567	CYS	2.4
1	A	361	CYS	2.4
1	A	30	TYR	2.4
1	A	506	THR	2.4
1	A	574	LYS	2.4
1	A	355	THR	2.3
1	A	558	CYS	2.3
1	A	362	ALA	2.3
1	A	433	VAL	2.3
1	A	530	VAL	2.3
1	A	94	GLN	2.2
1	A	550	ASP	2.2
1	A	368	GLU	2.2
1	A	77	VAL	2.2
1	A	364	ALA	2.2
1	A	523	ILE	2.2
1	A	563	ASP	2.2
1	A	176	ALA	2.2
1	A	555	VAL	2.2
1	A	177	CYS	2.2
1	A	418	VAL	2.1
1	A	516	LEU	2.1
1	A	505	GLU	2.1
1	A	435	SER	2.1
1	A	381	VAL	2.1
1	A	360	CYS	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	374	PHE	2.1
1	A	560	LYS	2.1
1	A	533	VAL	2.0
1	A	306	ALA	2.0
1	A	393	GLU	2.0
1	A	532	LEU	2.0
1	A	575	LEU	2.0
1	A	512	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.