



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 21, 2026 – 12:29 PM JST

PDB ID : 9UM6 / pdb_00009um6
Title : CaPETaseM9 SEC loop of 10CL+E289P variant
Authors : Kim, K.; Ki, D.; Park, J.
Deposited on : 2025-04-21
Resolution : 1.66 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.66 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.66-1.66)
Clashscore	190562	2662 (1.66-1.66)
Ramachandran outliers	187476	2621 (1.66-1.66)
Sidechain outliers	187428	2621 (1.66-1.66)
RSRZ outliers	180081	2564 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	267	<div> <div>4%</div> <div> <div></div> <div>91%</div> <div>6%</div> </div> <div></div> </div>
1	B	267	<div> <div>6%</div> <div> <div></div> <div>91%</div> <div>5%</div> </div> <div></div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4153 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cutinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	260	Total	C	N	O	S	0	0	0
			1981	1258	344	369	10			
1	B	259	Total	C	N	O	S	0	0	0
			1964	1247	341	366	10			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	39	GLU	ALA	engineered mutation	UNP A0A1M7II12
A	40	PRO	GLN	engineered mutation	UNP A0A1M7II12
A	109	ALA	ASN	engineered mutation	UNP A0A1M7II12
A	129	THR	VAL	engineered mutation	UNP A0A1M7II12
A	155	ARG	ALA	engineered mutation	UNP A0A1M7II12
A	180	CYS	LEU	engineered mutation	UNP A0A1M7II12
A	196	THR	GLY	engineered mutation	UNP A0A1M7II12
A	198	LYS	ARG	engineered mutation	UNP A0A1M7II12
A	202	CYS	ALA	engineered mutation	UNP A0A1M7II12
A	242	CYS	ARG	engineered mutation	UNP A0A1M7II12
A	283	ARG	SER	engineered mutation	UNP A0A1M7II12
A	284	ASP	THR	engineered mutation	UNP A0A1M7II12
A	288	GLY	ALA	engineered mutation	UNP A0A1M7II12
A	291	CYS	SER	engineered mutation	UNP A0A1M7II12
A	300	LEU	-	expression tag	UNP A0A1M7II12
A	301	GLU	-	expression tag	UNP A0A1M7II12
A	302	HIS	-	expression tag	UNP A0A1M7II12
A	303	HIS	-	expression tag	UNP A0A1M7II12
A	304	HIS	-	expression tag	UNP A0A1M7II12
A	305	HIS	-	expression tag	UNP A0A1M7II12
B	39	GLU	ALA	engineered mutation	UNP A0A1M7II12
B	40	PRO	GLN	engineered mutation	UNP A0A1M7II12
B	109	ALA	ASN	engineered mutation	UNP A0A1M7II12
B	129	THR	VAL	engineered mutation	UNP A0A1M7II12
B	155	ARG	ALA	engineered mutation	UNP A0A1M7II12

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	180	CYS	LEU	engineered mutation	UNP A0A1M7II12
B	196	THR	GLY	engineered mutation	UNP A0A1M7II12
B	198	LYS	ARG	engineered mutation	UNP A0A1M7II12
B	202	CYS	ALA	engineered mutation	UNP A0A1M7II12
B	242	CYS	ARG	engineered mutation	UNP A0A1M7II12
B	283	ARG	SER	engineered mutation	UNP A0A1M7II12
B	284	ASP	THR	engineered mutation	UNP A0A1M7II12
B	288	GLY	ALA	engineered mutation	UNP A0A1M7II12
B	291	CYS	SER	engineered mutation	UNP A0A1M7II12
B	300	LEU	-	expression tag	UNP A0A1M7II12
B	301	GLU	-	expression tag	UNP A0A1M7II12
B	302	HIS	-	expression tag	UNP A0A1M7II12
B	303	HIS	-	expression tag	UNP A0A1M7II12
B	304	HIS	-	expression tag	UNP A0A1M7II12
B	305	HIS	-	expression tag	UNP A0A1M7II12

- Molecule 2 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	126	Total O 126 126	0	0
2	B	82	Total O 82 82	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cutinase

Chain A: 



- Molecule 1: Cutinase

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	41.13Å 116.36Å 142.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.08 – 1.66 34.08 – 1.66	Depositor EDS
% Data completeness (in resolution range)	99.8 (34.08-1.66) 99.8 (34.08-1.66)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.23 (at 1.66Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
R, R_{free}	0.185 , 0.209 0.198 , 0.218	Depositor DCC
R_{free} test set	4060 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	18.2	Xtriage
Anisotropy	0.033	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 27.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4153	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	1/2041 (0.0%)	1.22	7/2792 (0.3%)
1	B	0.85	2/2023 (0.1%)	1.25	5/2768 (0.2%)
All	All	0.86	3/4064 (0.1%)	1.24	12/5560 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
All	All	0	4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	102	ILE	C-N	10.20	1.47	1.33
1	B	101	PHE	C-N	7.44	1.43	1.33
1	A	248	PHE	C-O	-5.87	1.19	1.24

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	102	ILE	O-C-N	13.15	139.01	122.57
1	B	101	PHE	CA-C-O	-6.63	114.74	122.37
1	A	63	PRO	N-CA-C	6.31	121.79	114.03
1	B	260	PHE	CA-CB-CG	6.00	119.80	113.80
1	A	260	PHE	CA-CB-CG	5.52	119.32	113.80
1	A	99	PRO	O-C-N	5.50	129.69	122.86
1	B	248	PHE	CA-CB-CG	5.48	119.28	113.80
1	A	64	PHE	CA-CB-CG	-5.41	108.39	113.80
1	A	44	ASN	CB-CA-C	5.22	115.34	110.17

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	101	PHE	CA-CB-CG	5.15	118.95	113.80
1	A	160	ARG	CD-NE-CZ	5.05	131.47	124.40
1	B	284	ASP	CA-CB-CG	5.02	117.62	112.60

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	137	ARG	Sidechain
1	A	176	ARG	Sidechain
1	B	137	ARG	Sidechain
1	B	182	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1981	0	1899	6	0
1	B	1964	0	1885	6	0
2	A	126	0	0	0	0
2	B	82	0	0	0	0
All	All	4153	0	3784	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:102:ILE:HA	1:A:130:ILE:HG12	1.93	0.50
1:B:98:VAL:HG12	1:B:99:PRO:O	2.12	0.49
1:A:165:ALA:O	1:A:188:ILE:HA	2.16	0.46
1:B:155:ARG:HH11	1:B:155:ARG:HG2	1.80	0.46
1:B:165:ALA:O	1:B:188:ILE:HA	2.17	0.44
1:A:102:ILE:HD13	1:B:170:MET:HE3	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:102:ILE:HG13	1:B:102:ILE:O	2.18	0.43
1:A:102:ILE:HG21	1:B:170:MET:CE	2.49	0.43
1:A:248:PHE:N	1:A:249:PRO:CD	2.82	0.42
1:A:251:SER:O	1:A:252:ALA:C	2.64	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	258/267 (97%)	253 (98%)	5 (2%)	0	100	100
1	B	257/267 (96%)	251 (98%)	6 (2%)	0	100	100
All	All	515/534 (96%)	504 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	207/212 (98%)	207 (100%)	0	100	100
1	B	204/212 (96%)	202 (99%)	2 (1%)	68	52
All	All	411/424 (97%)	409 (100%)	2 (0%)	81	72

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	197	GLU
1	B	291	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	276	GLN
1	A	302	HIS
1	B	181	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	260/267 (97%)	-0.01	11 (4%) 40 44	9, 16, 26, 66	0
1	B	259/267 (97%)	0.49	17 (6%) 24 27	13, 22, 39, 63	0
All	All	519/534 (97%)	0.24	28 (5%) 31 35	9, 19, 34, 66	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	300	LEU	5.6
1	A	302	HIS	5.3
1	B	153	PRO	5.2
1	B	101	PHE	5.1
1	B	102	ILE	5.0
1	B	41	ALA	4.7
1	A	101	PHE	3.9
1	B	151	ARG	3.7
1	A	301	GLU	3.7
1	B	72	VAL	3.7
1	A	43	ASP	3.5
1	A	100	GLY	3.3
1	A	151	ARG	3.2
1	B	283	ARG	3.1
1	B	154	VAL	3.1
1	B	100	GLY	2.9
1	B	66	VAL	2.8
1	B	150	THR	2.7
1	A	286	LEU	2.6
1	A	102	ILE	2.5
1	B	299	PHE	2.4
1	A	287	PHE	2.3
1	B	286	LEU	2.3
1	B	73	GLY	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	152	SER	2.2
1	A	202	CYS	2.1
1	B	291	CYS	2.1
1	B	75	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.