



Full wwPDB EM Validation Report ⓘ

Apr 5, 2026 – 09:00 PM UTC

PDB ID : 9UP2 / pdb_00009up2
EMDB ID : EMD-64387
Title : PvdL-E2-C3-A3-PCP3 in complex with MLP (NRPS cross-module)
Authors : Cao, W.; Wang, J.; Wang, Z.
Deposited on : 2025-04-27
Resolution : 3.33 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY



The reported resolution of this entry is 3.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	14484 (2.83 - 3.83)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1536	 97%
2	B	61	 98%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 25216 atoms, of which 12496 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PvdL.

Mol	Chain	Residues	Atoms							AltConf	Trace
1	A	1536	Total	C	H	N	O	P	S	0	0
			24143	7701	11978	2181	2244	1	38		

- Molecule 2 is a protein called MbtH-like domain-containing protein.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
2	B	61	996	325	487	87	94	3	0	0


- Molecule 3 is 2,4-DIAMINO BUTYRIC ACID (CCD ID: DAB) (formula: $C_4H_{10}N_2O_2$).



Mol	Chain	Residues	Atoms					AltConf
			Total	C	H	N	O	
3	A	1	17	4	9	2	2	0
3	A	1	17	4	10	2	1	0

- Molecule 4 is ADENOSINE-5'-TRIPHOSPHATE (CCD ID: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$)

(labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					AltConf	
4	A	1	Total	C	H	N	O	P	0
			43	10	12	5	13	3	


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PvdL



• Molecule 2: MbtH-like domain-containing protein



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	165245	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING ONLY	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	64000	Depositor
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.042	Depositor
Minimum map value	-0.012	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.001	Depositor
Recommended contour level	0.004	Depositor
Map size (\AA)	330.0, 330.0, 330.0	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.1, 1.1, 1.1	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 4HH, ATP, DAB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.79	0/12425	1.34	27/16901 (0.2%)
2	B	0.73	0/523	1.30	0/708
All	All	0.79	0/12948	1.34	27/17609 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4

There are no bond length outliers.

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	4016	ASP	CA-CB-CG	7.95	120.55	112.60
1	A	3994	ARG	NE-CZ-NH2	6.71	125.24	119.20
1	A	3057	ARG	NE-CZ-NH2	6.57	125.11	119.20
1	A	3649	HIS	CA-CB-CG	6.17	119.97	113.80
1	A	3272	ILE	N-CA-C	6.11	117.58	110.62
1	A	4128	ARG	NE-CZ-NH2	5.99	124.59	119.20
1	A	3011	ARG	NE-CZ-NH2	5.96	124.56	119.20
1	A	3821	ARG	NE-CZ-NH2	5.86	124.48	119.20
1	A	4289	ARG	NE-CZ-NH2	5.82	124.44	119.20
1	A	3278	ARG	NE-CZ-NH2	5.81	124.43	119.20
1	A	3057	ARG	CD-NE-CZ	5.78	132.49	124.40
1	A	3437	ARG	NE-CZ-NH2	5.62	124.26	119.20
1	A	3411	ARG	NE-CZ-NH2	5.47	124.13	119.20
1	A	3914	PHE	CA-CB-CG	5.36	119.16	113.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	3393	ARG	NE-CZ-NH2	5.35	124.02	119.20
1	A	2908	ARG	NE-CZ-NH2	5.30	123.97	119.20
1	A	3134	ASN	N-CA-C	5.25	116.82	108.79
1	A	3930	ARG	NE-CZ-NH2	5.24	123.92	119.20
1	A	3449	ARG	NE-CZ-NH2	5.22	123.90	119.20
1	A	3617	ARG	NE-CZ-NH2	5.22	123.90	119.20
1	A	3486	ARG	NE-CZ-NH2	5.14	123.83	119.20
1	A	4256	ARG	NE-CZ-NH2	5.14	123.82	119.20
1	A	3271	GLY	CA-C-N	5.12	127.76	120.53
1	A	3271	GLY	C-N-CA	5.12	127.76	120.53
1	A	3141	GLN	OE1-CD-NE2	-5.08	117.52	122.60
1	A	3013	ARG	NE-CZ-NH2	5.02	123.72	119.20
1	A	4301	ARG	NE-CZ-NH2	5.01	123.71	119.20

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	2885	ARG	Sidechain
1	A	3475	ARG	Sidechain
1	A	3790	TYR	Sidechain
1	A	4083	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	12165	11978	11977	1	0
2	B	509	487	484	0	0
3	A	15	19	18	0	0
4	A	31	12	12	0	0
All	All	12720	12496	12491	1	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (1) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:4020:PHE:CE2	1:A:4039:THR:HG22	2.53	0.43

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1533/1536 (100%)	1479 (96%)	53 (4%)	1 (0%)	48	76
2	B	59/61 (97%)	58 (98%)	1 (2%)	0	100	100
All	All	1592/1597 (100%)	1537 (96%)	54 (3%)	1 (0%)	49	76

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2848	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1259/1259 (100%)	1244 (99%)	15 (1%)	63	74
2	B	54/55 (98%)	53 (98%)	1 (2%)	50	68
All	All	1313/1314 (100%)	1297 (99%)	16 (1%)	61	74

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2793	THR
1	A	2795	ILE
1	A	2797	HIS
1	A	2839	ASP
1	A	3057	ARG
1	A	3064	ILE
1	A	3121	GLN
1	A	3134	ASN
1	A	3511	ARG
1	A	3568	LEU
1	A	3862	GLN
1	A	3868	ILE
1	A	3911	SER
1	A	4016	ASP
1	A	4214	LEU
2	B	30	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2797	HIS
1	A	2811	ASN
1	A	2856	HIS
1	A	2915	HIS
1	A	3045	GLN
1	A	3153	GLN
1	A	3383	ASN
1	A	3385	HIS
1	A	3499	ASN
1	A	3503	GLN
1	A	3572	ASN
1	A	3587	GLN
1	A	3912	GLN
1	A	4279	HIS
2	B	11	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	4HH	A	4280	3,1	22,26,27	1.00	1 (4%)	27,35,37	1.43	2 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	4HH	A	4280	3,1	-	9/33/35/37	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	4280	4HH	CJ-CK	2.49	1.56	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	4280	4HH	OG-CB-CA	5.04	113.05	108.14
1	A	4280	4HH	P-OG-CB	3.17	139.52	121.35

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	4280	4HH	O3P-CJ-CK-CL1
1	A	4280	4HH	O3P-CJ-CK-CL2
1	A	4280	4HH	NN-CL3-CM-OM
1	A	4280	4HH	ON-CL3-CM-OM
1	A	4280	4HH	ON-CL3-CM-CK

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
1	A	4280	4HH	NN-CL3-CM-CK
1	A	4280	4HH	CA-CB-OG-P
1	A	4280	4HH	O3P-CJ-CK-CM
1	A	4280	4HH	NR-CS-CT-SU

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	DAB	A	4403	1	5,6,7	0.90	0	1,6,8	0.18	0
4	ATP	A	4402	-	32,33,33	1.27	4 (12%)	48,52,52	1.36	7 (14%)
3	DAB	A	4401	-	6,7,7	0.66	0	4,8,8	0.85	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	DAB	A	4403	1	-	0/4/5/7	-
4	ATP	A	4402	-	-	11/22/38/38	0/3/3/3
3	DAB	A	4401	-	-	1/7/7/7	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	4402	ATP	PA-O3A	-2.69	1.56	1.59
4	A	4402	ATP	PB-O3B	-2.66	1.56	1.59
4	A	4402	ATP	PB-O3A	-2.43	1.56	1.59
4	A	4402	ATP	C5-C4	-2.04	1.35	1.39

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	4402	ATP	C5-C4-N3	-3.21	122.30	126.72
4	A	4402	ATP	C4-C5-N7	3.15	114.18	110.58
4	A	4402	ATP	N3-C4-N9	2.52	131.46	127.17
4	A	4402	ATP	O2G-PG-O3B	2.35	112.50	104.64
4	A	4402	ATP	C6-C5-N7	-2.15	127.94	132.09
4	A	4402	ATP	C2'-C1'-N9	2.09	118.51	113.30
4	A	4402	ATP	C2-N1-C6	-2.01	115.44	118.73

There are no chirality outliers.

All (12) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	4402	ATP	PB-O3B-PG-O2G
4	A	4402	ATP	C4'-C5'-O5'-PA
4	A	4402	ATP	O4'-C4'-C5'-O5'
4	A	4402	ATP	C3'-C4'-C5'-O5'
4	A	4402	ATP	C2'-C1'-N9-C4
4	A	4402	ATP	PB-O3A-PA-O5'
4	A	4402	ATP	C5'-O5'-PA-O2A
4	A	4402	ATP	C5'-O5'-PA-O3A
3	A	4401	DAB	O-C-CA-N
4	A	4402	ATP	C2'-C1'-N9-C8
4	A	4402	ATP	O4'-C1'-N9-C8
4	A	4402	ATP	PB-O3B-PG-O1G

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-64387. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 150




Y Index: 150




Z Index: 150


6.2.2 Raw map



X Index: 150



Y Index: 150




Z Index: 150


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 182




Y Index: 139




Z Index: 142


6.3.2 Raw map



X Index: 156



Y Index: 139




Z Index: 142


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z


6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.004. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 238 nm^3 ; this corresponds to an approximate mass of 215 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.300 \AA^{-1}

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.300 \AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.33	-	-
Author-provided FSC curve	3.31	3.65	3.34
Unmasked-calculated*	3.65	4.14	3.70

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-64387 and PDB model 9UP2. Per-residue inclusion information can be found in [section 3](#) on [page 5](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.004 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.004).

9.4 Atom inclusion ⓘ



At the recommended contour level, 97% of all backbone atoms, 97% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.004) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.9660	<div></div> 0.4250
A	<div></div> 0.9720	<div></div> 0.4220
B	<div></div> 0.9860	<div></div> 0.4980

