



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 07:12 PM UTC

PDB ID : 9VLA / pdb_00009vla
Title : Crystal structure of PNGase Bb in complex with MUXF glycopeptide
Authors : Wu, T.T.; Voglmeir, J.; Wei, B.Q.
Deposited on : 2025-06-24
Resolution : 2.16 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	561	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> % 87% 9% ... </div> </div>
1	B	561	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> % 85% 10% ... </div> </div>
2	H	3	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, yellow, green);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> 100% </div> </div>
2	I	3	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green, yellow);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> 67% 33% </div> </div>
3	C	6	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green, yellow);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> 50% 50% </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	E	6	<div><div></div><div>17%</div><div>83%</div></div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8550 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PNGase Bb.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	539	Total	C	N	O	S	0	0	0
			4051	2511	689	842	9			
1	B	539	Total	C	N	O	S	0	0	0
			4050	2512	688	840	10			

- Molecule 2 is a protein called ASN-DGL-DSN.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	H	3	Total	C	N	O	0	0	0
			24	12	4	8			
2	I	3	Total	C	N	O	0	0	0
			24	12	4	8			

- Molecule 3 is an oligosaccharide called beta-D-xylopyranose-(1-2)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-galactopyranose-(1-4)-[beta-L-fucopyranose-(1-3)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	6	Total	C	N	O	0	0	0
			69	39	2	28			
3	E	6	Total	C	N	O	0	0	0
			69	39	2	28			

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Cl	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	139	Total	O	0	0
			139	139		
5	B	118	Total	O	0	0
			118	118		
5	H	1	Total	O	0	0
			1	1		
5	I	3	Total	O	0	0
			3	3		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PNGase Bb



• Molecule 1: PNGase Bb



• Molecule 2: ASN-DGL-DSN



• Molecule 2: ASN-DGL-DSN



- Molecule 3: beta-D-xylopyranose-(1-2)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-galactopyranose-(1-4)-[beta-L-fucopyranose-(1-3)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:  50% 50%

NAG1	NAG2	BMA3	XYP4	MAN5	FUL6
------	------	------	------	------	------

- Molecule 3: beta-D-xylopyranose-(1-2)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-galactopyranose-(1-4)-[beta-L-fucopyranose-(1-3)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  17% 83%

NAG1	NAG2	BMA3	XYP4	MAN5	FUL6
------	------	------	------	------	------

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	136.72Å 97.89Å 96.32Å 90.00° 122.02° 90.00°	Depositor
Resolution (Å)	33.47 – 2.16 33.47 – 2.16	Depositor EDS
% Data completeness (in resolution range)	99.2 (33.47-2.16) 99.3 (33.47-2.16)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.65 (at 2.16Å)	Xtriage
Refinement program	PHENIX (1.19_4092: ???)	Depositor
R, R_{free}	0.208 , 0.261 0.211 , 0.263	Depositor DCC
R_{free} test set	2000 reflections (3.46%)	wwPDB-VP
Wilson B-factor (Å ²)	44.8	Xtriage
Anisotropy	0.240	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 46.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8550	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, NGA, DSN, BMA, FUL, MAN, XYP, DGL, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.48	0/4139	0.66	0/5662
1	B	0.53	0/4138	0.72	1/5660 (0.0%)
2	H	0.79	0/7	1.10	0/8
2	I	0.75	0/7	0.74	0/8
All	All	0.50	0/8291	0.69	1/11338 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
2	H	0	1
2	I	0	1
All	All	0	5

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	524	ASP	CA-CB-CG	5.26	117.86	112.60

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	277	ARG	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	B	288	ARG	Sidechain
1	B	436	ARG	Sidechain
2	H	2	DGL	Peptide
2	I	2	DGL	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4051	0	3849	32	0
1	B	4050	0	3849	32	0
2	H	24	0	19	3	0
2	I	24	0	18	0	0
3	C	69	0	51	0	0
3	E	69	0	51	1	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	139	0	0	4	0
5	B	118	0	0	4	0
5	H	1	0	0	1	0
5	I	3	0	0	0	0
All	All	8550	0	7837	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:208:THR:HB	1:A:323:THR:HG22	1.73	0.69
1:A:375:THR:HG23	1:A:392:THR:HG22	1.74	0.68
1:A:456:MET:HA	1:A:456:MET:HE2	1.79	0.64
3:E:3:BMA:H4	3:E:5:MAN:H3	1.81	0.63
1:B:369:ASN:ND2	1:B:398:GLN:HE21	1.96	0.63
1:B:201:THR:HG22	1:B:202:GLN:HG3	1.81	0.62
1:A:105:GLN:HE21	1:A:106:GLU:H	1.48	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:213:GLY:O	1:B:216:GLU:HG2	1.99	0.61
1:A:233:ALA:HB3	5:A:770:HOH:O	2.01	0.59
1:B:369:ASN:HD22	1:B:398:GLN:HE21	1.52	0.58
1:B:528:THR:HG22	1:B:548:THR:OG1	2.04	0.57
1:A:494:ASN:HB2	5:A:778:HOH:O	2.05	0.55
1:B:263:PHE:CD1	1:B:264:PRO:HD2	2.42	0.55
1:B:528:THR:HB	1:B:549:SER:HB3	1.88	0.55
1:A:38:VAL:HG11	1:A:60:ILE:HD13	1.89	0.55
1:B:334:LEU:HG	1:B:383:SER:HB3	1.88	0.55
1:A:232:ASP:O	1:A:236:THR:N	2.38	0.53
1:A:44:PHE:CZ	1:A:54:GLN:HG2	2.45	0.52
1:A:119:LYS:NZ	5:A:708:HOH:O	2.41	0.52
1:A:204:GLN:HE22	1:A:485:LEU:HD22	1.73	0.51
1:A:461:TYR:CE2	2:H:3:DSN:HB3	2.46	0.51
1:B:427:SER:HB2	1:B:436:ARG:HG2	1.93	0.51
1:B:204:GLN:NE2	1:B:485:LEU:HD22	2.27	0.50
1:A:231:SER:O	1:A:232:ASP:C	2.55	0.49
1:B:204:GLN:HE22	1:B:485:LEU:HD22	1.77	0.49
1:A:232:ASP:O	1:A:233:ALA:C	2.56	0.48
1:A:177:VAL:O	5:A:701:HOH:O	2.20	0.48
1:B:539:LYS:NZ	5:B:705:HOH:O	2.39	0.48
1:B:217:GLN:NE2	5:B:702:HOH:O	2.28	0.48
1:A:427:SER:HB2	1:A:436:ARG:HG2	1.96	0.48
1:B:233:ALA:O	1:B:237:GLN:HB2	2.14	0.47
1:A:202:GLN:HB2	1:A:329:THR:HB	1.97	0.47
1:B:238:MET:HE3	1:B:455:PHE:HB2	1.96	0.47
1:B:175:VAL:HG11	1:B:194:PRO:HG3	1.96	0.47
1:B:87:ASP:OD1	1:B:87:ASP:N	2.49	0.46
1:B:173:GLN:HB2	1:B:328:LEU:HB2	1.97	0.45
1:A:475:ARG:HD2	1:A:487:PRO:O	2.16	0.45
1:A:65:GLU:CD	1:A:65:GLU:H	2.24	0.45
1:A:455:PHE:CE1	1:A:456:MET:HE3	2.52	0.45
1:B:341:LEU:HD12	1:B:377:SER:O	2.17	0.45
1:A:445:PRO:HG2	1:A:470:SER:HB2	1.98	0.44
1:A:405:VAL:HG12	1:A:406:SER:O	2.17	0.44
1:A:188:SER:HB2	1:A:312:SER:HB2	1.99	0.44
1:A:233:ALA:HB1	1:A:456:MET:SD	2.57	0.43
1:A:257:ALA:O	1:A:376:ILE:HD11	2.19	0.43
1:B:231:SER:O	1:B:233:ALA:N	2.51	0.43
1:A:232:ASP:OD1	1:A:232:ASP:N	2.51	0.43
1:A:185:LYS:HB3	1:A:185:LYS:HE2	1.80	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:97:VAL:HB	1:B:127:LEU:HD11	2.00	0.43
1:A:308:ARG:CZ	1:A:308:ARG:HB2	2.47	0.43
1:B:314:ASN:ND2	5:B:701:HOH:O	2.21	0.43
1:A:461:TYR:CD2	2:H:3:DSN:HB3	2.54	0.42
1:B:81:VAL:HG13	1:B:107:PRO:HB3	2.01	0.42
1:B:191:LEU:HD12	1:B:309:LEU:O	2.19	0.42
1:A:474:LYS:HB2	1:A:474:LYS:NZ	2.34	0.42
1:B:402:THR:HG22	1:B:414:SER:HB3	2.00	0.42
1:B:238:MET:HG3	1:B:455:PHE:CD2	2.55	0.42
1:B:14:ILE:N	5:B:713:HOH:O	2.53	0.41
1:B:62:PRO:HG3	1:B:128:PHE:O	2.21	0.41
1:B:200:LEU:HD21	1:B:328:LEU:HD22	2.02	0.41
1:A:89:THR:OG1	1:A:210:LYS:HE3	2.21	0.41
1:B:111:THR:HA	1:B:112:PRO:HD3	1.95	0.41
1:B:253:ASP:OD1	1:B:308:ARG:N	2.38	0.41
1:A:350:SER:O	1:A:370:GLY:HA2	2.21	0.40
1:B:89:THR:OG1	1:B:210:LYS:HE3	2.22	0.40
2:H:1:ASN:OD1	5:H:101:HOH:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	537/561 (96%)	516 (96%)	17 (3%)	4 (1%)	18	13
1	B	537/561 (96%)	515 (96%)	18 (3%)	4 (1%)	18	13
All	All	1074/1122 (96%)	1031 (96%)	35 (3%)	8 (1%)	18	13

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	232	ASP
1	A	231	SER
1	A	233	ALA
1	B	233	ALA
1	A	232	ASP
1	B	237	GLN
1	B	271	ILE
1	A	271	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	444/462 (96%)	436 (98%)	8 (2%)	51	58
1	B	443/462 (96%)	429 (97%)	14 (3%)	34	36
2	H	1/1 (100%)	1 (100%)	0	100	100
2	I	1/1 (100%)	1 (100%)	0	100	100
All	All	889/926 (96%)	867 (98%)	22 (2%)	42	45

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	214	CYS
1	A	231	SER
1	A	234	VAL
1	A	237	GLN
1	A	311	LEU
1	A	323	THR
1	A	354	THR
1	A	450	LEU
1	B	61	THR
1	B	163	GLU
1	B	191	LEU
1	B	195	MET
1	B	201	THR
1	B	214	CYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	232	ASP
1	B	237	GLN
1	B	309	LEU
1	B	321	ASN
1	B	350	SER
1	B	415	MET
1	B	431	LYS
1	B	450	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	54	GLN
1	A	105	GLN
1	A	204	GLN
1	A	340	ASN
1	A	360	ASN
1	A	409	ASN
1	A	424	GLN
1	A	472	ASN
1	A	514	HIS
1	B	43	HIS
1	B	202	GLN
1	B	343	GLN
1	B	369	ASN
1	B	400	ASN
1	B	424	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	C	1	3,2	14,14,15	0.48	0	17,19,21	2.11	6 (35%)
3	NGA	C	2	3	14,14,15	0.57	0	17,19,21	2.06	4 (23%)
3	BMA	C	3	3	11,11,12	0.60	0	15,15,17	0.69	0
3	XYP	C	4	3	9,9,10	0.31	0	10,12,14	0.83	0
3	MAN	C	5	3	11,11,12	0.35	0	15,15,17	0.64	0
3	FUL	C	6	3	10,10,11	0.60	0	14,14,16	1.72	2 (14%)
3	NAG	E	1	3,2	14,14,15	0.40	0	17,19,21	1.70	3 (17%)
3	NGA	E	2	3	14,14,15	0.40	0	17,19,21	1.08	2 (11%)
3	BMA	E	3	3	11,11,12	0.56	0	15,15,17	0.60	0
3	XYP	E	4	3	9,9,10	0.19	0	10,12,14	0.33	0
3	MAN	E	5	3	11,11,12	0.41	0	15,15,17	0.30	0
3	FUL	E	6	3	10,10,11	0.56	0	14,14,16	1.47	2 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	3,2	-	2/6/23/26	0/1/1/1
3	NGA	C	2	3	-	2/6/23/26	0/1/1/1
3	BMA	C	3	3	-	1/2/19/22	0/1/1/1
3	XYP	C	4	3	-	-	0/1/1/1
3	MAN	C	5	3	-	2/2/19/22	1/1/1/1
3	FUL	C	6	3	-	-	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	E	1	3,2	-	2/6/23/26	0/1/1/1
3	NGA	E	2	3	-	0/6/23/26	0/1/1/1
3	BMA	E	3	3	-	2/2/19/22	0/1/1/1
3	XYP	E	4	3	-	-	0/1/1/1
3	MAN	E	5	3	-	2/2/19/22	1/1/1/1
3	FUL	E	6	3	-	-	0/1/1/1

There are no bond length outliers.

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	2	NGA	C4-C3-C2	6.47	120.50	111.02
3	C	6	FUL	C1-C2-C3	-4.91	102.50	109.64
3	C	1	NAG	O3-C3-C4	4.50	120.98	110.38
3	E	1	NAG	O3-C3-C4	4.24	120.38	110.38
3	C	1	NAG	O5-C1-C2	-4.23	104.75	111.29
3	C	1	NAG	O3-C3-C2	-3.74	101.64	109.40
3	E	1	NAG	O5-C1-C2	-3.64	105.66	111.29
3	C	2	NGA	C2-N2-C7	3.54	127.65	122.90
3	E	6	FUL	C1-C2-C3	-3.40	104.69	109.64
3	E	1	NAG	O3-C3-C2	-3.07	103.03	109.40
3	C	1	NAG	C4-C3-C2	-2.97	106.67	111.02
3	C	2	NGA	O5-C1-C2	-2.74	107.06	111.29
3	E	6	FUL	O2-C2-C3	2.55	115.44	110.15
3	E	2	NGA	C1-C2-N2	2.38	114.18	110.43
3	C	1	NAG	C3-C4-C5	-2.29	106.08	110.23
3	E	2	NGA	O4-C4-C3	2.24	115.65	110.38
3	C	2	NGA	O4-C4-C3	2.18	115.52	110.38
3	C	1	NAG	C1-C2-N2	2.06	113.68	110.43
3	C	6	FUL	O2-C2-C3	2.05	114.40	110.15

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	1	NAG	C8-C7-N2-C2
3	C	1	NAG	O7-C7-N2-C2
3	E	1	NAG	C8-C7-N2-C2
3	C	2	NGA	O5-C5-C6-O6
3	E	5	MAN	O5-C5-C6-O6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	E	5	MAN	C4-C5-C6-O6
3	C	5	MAN	O5-C5-C6-O6
3	E	1	NAG	O7-C7-N2-C2
3	C	2	NGA	C4-C5-C6-O6
3	C	5	MAN	C4-C5-C6-O6
3	E	3	BMA	C4-C5-C6-O6
3	E	3	BMA	O5-C5-C6-O6
3	C	3	BMA	C4-C5-C6-O6


All (2) ring outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	E	5	MAN	C1-C2-C3-C4-C5-O5
3	C	5	MAN	C1-C2-C3-C4-C5-O5

2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	3	BMA	1	0
3	E	5	MAN	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	539/561 (96%)	0.16	6 (1%) 78 80	26, 46, 76, 117	0
1	B	539/561 (96%)	0.32	5 (0%) 81 83	28, 49, 82, 120	0
2	H	1/3 (33%)	-0.25	0 100 100	47, 47, 47, 47	0
2	I	1/3 (33%)	1.24	0 100 100	54, 54, 54, 54	0
All	All	1080/1128 (95%)	0.24	11 (1%) 79 82	26, 48, 79, 120	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	216	GLU	3.5
1	B	216	GLU	3.3
1	B	234	VAL	2.9
1	B	63	PRO	2.4
1	A	459	ASN	2.3
1	A	306	LYS	2.2
1	B	459	ASN	2.2
1	A	66	CYS	2.2
1	A	144	ASP	2.2
1	B	232	ASP	2.1
1	A	225	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	DGL	I	2	9/10	0.75	0.17	46,56,61,64	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	DSN	H	3	7/7	0.76	0.14	57,61,76,106	0
2	DSN	I	3	7/7	0.76	0.12	63,67,81,83	0
2	DGL	H	2	9/10	0.78	0.16	41,49,56,57	0

6.3 Carbohydrates




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NAG	C	1	14/15	-	-	35,46,52,52	0
3	NGA	C	2	14/15	-	-	41,49,56,56	0
3	BMA	C	3	11/12	-	-	45,52,57,58	0
3	XYP	C	4	9/10	-	-	61,66,75,83	0
3	MAN	C	5	11/12	-	-	49,51,63,64	0
3	FUL	C	6	10/11	-	-	38,46,49,56	0
3	XYP	E	4	9/10	0.68	0.13	118,124,128,129	0
3	MAN	E	5	11/12	0.84	0.10	120,130,135,140	0
3	FUL	E	6	10/11	0.86	0.12	52,56,59,61	0
3	BMA	E	3	11/12	0.90	0.07	103,112,117,126	0
3	NAG	E	1	14/15	0.93	0.09	48,56,59,63	0
3	NGA	E	2	14/15	0.93	0.09	58,68,88,98	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	CL	A	601	1/1	0.93	0.12	30,30,30,30	0
4	CL	B	601	1/1	0.93	0.22	30,30,30,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.