



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 4, 2026 – 10:09 PM UTC

PDB ID : 9WHA / pdb_00009wha
Title : Crystal Structure of Bovine STING bound to 2'3'-cGAMP
Authors : Zhao, Z.Z.; Li, X.
Deposited on : 2025-08-26
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	195	<div> <div>17%</div> <div>85%</div> <div>11%</div> <div>.</div> </div>
1	B	195	<div> <div>16%</div> <div>88%</div> <div>7%</div> <div>5%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3346 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Stimulator of interferon genes protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	187	Total	C	N	O	S	0	0	0
			1490	938	268	279	5			
1	B	186	Total	C	N	O	S	0	0	0
			1482	932	267	278	5			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	154	MET	-	initiating methionine	UNP Q2KI99
A	341	LEU	-	expression tag	UNP Q2KI99
A	342	GLU	-	expression tag	UNP Q2KI99
A	343	HIS	-	expression tag	UNP Q2KI99
A	344	HIS	-	expression tag	UNP Q2KI99
A	345	HIS	-	expression tag	UNP Q2KI99
A	346	HIS	-	expression tag	UNP Q2KI99
A	347	HIS	-	expression tag	UNP Q2KI99
A	348	HIS	-	expression tag	UNP Q2KI99
B	154	MET	-	initiating methionine	UNP Q2KI99
B	341	LEU	-	expression tag	UNP Q2KI99
B	342	GLU	-	expression tag	UNP Q2KI99
B	343	HIS	-	expression tag	UNP Q2KI99
B	344	HIS	-	expression tag	UNP Q2KI99
B	345	HIS	-	expression tag	UNP Q2KI99
B	346	HIS	-	expression tag	UNP Q2KI99
B	347	HIS	-	expression tag	UNP Q2KI99
B	348	HIS	-	expression tag	UNP Q2KI99

- Molecule 2 is cGAMP (CCD ID: 1SY) (formula: $C_{20}H_{24}N_{10}O_{13}P_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 90	C 40	N 20	O 26	P 4	0	1


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	159	Total O 159 159	0	0
3	B	125	Total O 125 125	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Stimulator of interferon genes protein



- Molecule 1: Stimulator of interferon genes protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	50.73Å 57.91Å 132.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	53.05 – 1.60 53.05 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.3 (53.05-1.60) 99.2 (53.05-1.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.79 (at 1.60Å)	Xtriage
Refinement program	PHENIX (1.17.1_3660: ???)	Depositor
R, R_{free}	0.216 , 0.241 0.216 , 0.242	Depositor DCC
R_{free} test set	2572 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	16.1	Xtriage
Anisotropy	1.476	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 44.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3346	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 1SY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.15	0/1521	0.36	0/2063
1	B	0.13	0/1513	0.33	0/2052
All	All	0.14	0/3034	0.34	0/4115

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1490	0	1457	13	0
1	B	1482	0	1446	9	0
2	A	90	0	44	6	0
3	A	159	0	0	10	0
3	B	125	0	0	5	0
All	All	3346	0	2947	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:401[B]:1SY:O2'	1:B:263:THR:OG1	2.00	0.77
1:B:252:GLN:NE2	3:B:402:HOH:O	2.18	0.76
1:A:155:VAL:N	3:A:506:HOH:O	2.23	0.71
1:A:188:ASN:ND2	1:A:191:GLN:OE1	2.25	0.70
2:A:401[B]:1SY:N42	3:A:502:HOH:O	1.99	0.70
2:A:401[A]:1SY:H21	2:A:401[A]:1SY:O26	1.92	0.69
1:B:301:ASN:OD1	3:B:401:HOH:O	2.09	0.69
1:A:253:ARG:NH1	3:A:501:HOH:O	1.99	0.67
2:A:401[B]:1SY:H21	2:A:401[B]:1SY:O26	1.95	0.65
1:A:305:SER:OG	3:A:504:HOH:O	2.12	0.65
1:A:232:ARG:NH1	3:A:507:HOH:O	2.28	0.64
1:A:224:GLU:OE2	3:A:505:HOH:O	2.17	0.58
1:B:186:HIS:O	1:B:189:THR:OG1	2.22	0.57
1:A:246:GLU:OE2	1:A:253:ARG:NH2	2.39	0.53
1:B:316:GLU:O	3:B:403:HOH:O	2.19	0.52
1:B:307:ASN:ND2	3:B:408:HOH:O	2.41	0.51
1:A:179:ILE:HG23	1:A:190:LEU:HD23	1.94	0.50
1:A:181:ILE:HD12	3:A:640:HOH:O	2.11	0.50
1:B:306:GLN:NE2	3:B:411:HOH:O	2.46	0.48
1:A:173:PRO:HG2	3:A:622:HOH:O	2.13	0.48
1:B:334:ARG:O	1:B:338:ARG:HG3	2.16	0.45
1:A:232:ARG:HD2	3:A:592:HOH:O	2.17	0.45
1:A:299:LEU:O	1:A:306:GLN:HG2	2.16	0.45
1:A:227:GLN:HG2	1:A:241:THR:HG22	2.00	0.42
2:A:401[B]:1SY:C40	3:A:502:HOH:O	2.59	0.42
2:A:401[B]:1SY:O4'	1:B:167:TYR:HB2	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	185/195 (95%)	179 (97%)	4 (2%)	2 (1%)	11 2

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	184/195 (94%)	182 (99%)	2 (1%)	0	100	100
All	All	369/390 (95%)	361 (98%)	6 (2%)	2 (0%)	24	10

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	317	PRO
1	A	320	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	159/167 (95%)	156 (98%)	3 (2%)	50	27
1	B	158/167 (95%)	155 (98%)	3 (2%)	50	27
All	All	317/334 (95%)	311 (98%)	6 (2%)	50	27

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	180	GLN
1	A	339	GLU
1	A	341	LEU
1	B	190	LEU
1	B	315	GLN
1	B	337	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	273	GLN
1	B	157	HIS
1	B	184	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	196	HIS
1	B	306	GLN
1	B	332	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	1SY	A	401[B]	-	51,51,51	2.27	20 (39%)	70,80,80	2.04	14 (20%)
2	1SY	A	401[A]	-	51,51,51	2.28	20 (39%)	70,80,80	2.01	15 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1SY	A	401[B]	-	-	5/30/62/62	0/6/7/7

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1SY	A	401[A]	-	-	4/30/62/62	0/6/7/7

All (40) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401[B]	1SY	C36-N42	6.84	1.49	1.37
2	A	401[A]	1SY	C36-N42	6.52	1.48	1.37
2	A	401[A]	1SY	C36-N33	-5.61	1.31	1.37
2	A	401[B]	1SY	C36-N33	-4.56	1.32	1.37
2	A	401[A]	1SY	C6-N01	4.55	1.45	1.34
2	A	401[B]	1SY	C40-N41	4.52	1.44	1.34
2	A	401[B]	1SY	C6-N01	4.48	1.45	1.34
2	A	401[A]	1SY	C40-N41	4.42	1.44	1.34
2	A	401[A]	1SY	C40-N39	4.41	1.43	1.33
2	A	401[B]	1SY	C40-N39	4.39	1.43	1.33
2	A	401[A]	1SY	C40-N42	3.26	1.45	1.37
2	A	401[B]	1SY	C40-N42	3.22	1.45	1.37
2	A	401[A]	1SY	C38-C37	3.15	1.54	1.45
2	A	401[B]	1SY	C38-C37	3.12	1.54	1.45
2	A	401[B]	1SY	C5-C4	-3.03	1.33	1.39
2	A	401[B]	1SY	O43-C38	-2.96	1.18	1.23
2	A	401[A]	1SY	C5-C4	-2.89	1.33	1.39
2	A	401[A]	1SY	C38-N39	2.80	1.42	1.38
2	A	401[B]	1SY	C2'-C3'	-2.76	1.47	1.53
2	A	401[B]	1SY	C38-N39	2.71	1.42	1.38
2	A	401[B]	1SY	C37-N35	-2.70	1.33	1.39
2	A	401[A]	1SY	C37-N35	-2.69	1.33	1.39
2	A	401[A]	1SY	C3'-C4'	-2.66	1.46	1.52
2	A	401[B]	1SY	P27-O28	2.61	1.67	1.59
2	A	401[B]	1SY	C21-C32	-2.58	1.46	1.53
2	A	401[A]	1SY	C21-C32	-2.56	1.46	1.53
2	A	401[B]	1SY	C3'-C4'	-2.55	1.46	1.52
2	A	401[A]	1SY	P18-O20	2.52	1.66	1.59
2	A	401[A]	1SY	O43-C38	-2.50	1.19	1.23
2	A	401[A]	1SY	C2'-C3'	-2.49	1.47	1.53
2	A	401[A]	1SY	P27-O28	2.49	1.66	1.59
2	A	401[A]	1SY	C22-C21	-2.46	1.47	1.53
2	A	401[B]	1SY	P18-O20	2.45	1.66	1.59
2	A	401[B]	1SY	P18-O17	2.41	1.68	1.59
2	A	401[A]	1SY	P18-O17	2.34	1.68	1.59
2	A	401[A]	1SY	C5-N7	-2.33	1.34	1.39
2	A	401[B]	1SY	C22-C21	-2.30	1.47	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401[B]	1SY	C5-N7	-2.26	1.35	1.39
2	A	401[B]	1SY	C8-N9	-2.13	1.33	1.37
2	A	401[A]	1SY	C8-N9	-2.01	1.34	1.37

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401[A]	1SY	C37-C36-N33	6.44	109.23	106.25
2	A	401[B]	1SY	C32-N33-C34	-6.24	108.99	126.73
2	A	401[A]	1SY	C32-N33-C34	-5.99	109.71	126.73
2	A	401[A]	1SY	N3-C2-N1	-5.70	119.95	128.58
2	A	401[B]	1SY	N3-C2-N1	-5.70	119.96	128.58
2	A	401[B]	1SY	N9-C8-N7	-4.88	107.01	113.94
2	A	401[B]	1SY	C37-C36-N33	4.81	108.47	106.25
2	A	401[B]	1SY	C5-C4-N3	-4.77	120.14	126.72
2	A	401[A]	1SY	N9-C8-N7	-4.74	107.22	113.94
2	A	401[A]	1SY	C5-C4-N3	-4.53	120.48	126.72
2	A	401[B]	1SY	C2-N3-C4	3.47	120.30	111.83
2	A	401[A]	1SY	C2-N3-C4	3.42	120.18	111.83
2	A	401[B]	1SY	C5-N7-C8	3.34	108.70	103.45
2	A	401[B]	1SY	O30-P27-O28	3.28	120.03	106.70
2	A	401[A]	1SY	O30-P27-O28	3.26	119.98	106.70
2	A	401[B]	1SY	O4'-C1'-N9	3.21	114.25	108.09
2	A	401[A]	1SY	N33-C34-N35	-3.20	107.47	113.40
2	A	401[A]	1SY	C5-N7-C8	3.20	108.48	103.45
2	A	401[B]	1SY	N33-C34-N35	-2.93	107.97	113.40
2	A	401[A]	1SY	C4-C5-N7	-2.89	107.27	110.58
2	A	401[B]	1SY	C4-N9-C8	2.89	108.78	105.74
2	A	401[B]	1SY	C4-C5-N7	-2.82	107.36	110.58
2	A	401[B]	1SY	N3-C4-N9	2.67	131.70	127.17
2	A	401[A]	1SY	C4-N9-C8	2.59	108.46	105.74
2	A	401[A]	1SY	C5-C4-N9	2.53	108.57	105.81
2	A	401[A]	1SY	N3-C4-N9	2.22	130.95	127.17
2	A	401[A]	1SY	C34-N35-C37	2.16	108.12	104.26
2	A	401[B]	1SY	C5-C4-N9	2.15	108.16	105.81
2	A	401[A]	1SY	O4'-C1'-N9	2.15	112.21	108.09

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401[B]	1SY	C21-O20-P18-O17

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	401[A]	1SY	C4'-C3'-O28-P27
2	A	401[B]	1SY	C4'-C3'-O28-P27
2	A	401[A]	1SY	C2'-C3'-O28-P27
2	A	401[B]	1SY	C2'-C3'-O28-P27
2	A	401[A]	1SY	C21-O20-P18-O17
2	A	401[A]	1SY	C25-O26-P27-O30
2	A	401[B]	1SY	C25-O26-P27-O30
2	A	401[B]	1SY	C21-O20-P18-O44


There are no ring outliers.


2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401[B]	1SY	5	0
2	A	401[A]	1SY	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand 1SY A 401 (B)





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	187/195 (95%)	1.05	34 (18%) 3 3	15, 29, 65, 111	0
1	B	186/195 (95%)	1.09	32 (17%) 4 3	17, 31, 57, 75	0
All	All	373/390 (95%)	1.07	66 (17%) 4 3	15, 30, 64, 111	0

All (66) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	340	VAL	4.8
1	B	315	GLN	4.6
1	A	322	SER	4.6
1	B	316	GLU	4.5
1	A	341	LEU	4.5
1	A	320	GLY	4.4
1	A	318	ALA	4.3
1	A	192	GLY	3.9
1	A	282	GLU	3.9
1	A	191	GLN	3.8
1	A	278	GLY	3.7
1	A	307	ASN	3.7
1	A	317	PRO	3.6
1	B	303	PRO	3.6
1	B	317	PRO	3.5
1	A	277	ALA	3.4
1	B	155	VAL	3.4
1	A	193	ALA	3.4
1	A	340	VAL	3.3
1	A	337	GLU	3.2
1	A	270	ALA	3.2
1	B	322	SER	3.1
1	B	321	SER	3.1
1	B	230	ALA	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	279	PHE	3.0
1	B	193	ALA	3.0
1	B	282	GLU	3.0
1	A	321	SER	2.9
1	B	286	GLU	2.8
1	A	189	THR	2.8
1	B	320	GLY	2.8
1	A	281	ARG	2.7
1	B	338	ARG	2.7
1	A	304	GLU	2.6
1	A	274	ASP	2.6
1	A	303	PRO	2.6
1	B	276	ARG	2.6
1	B	307	ASN	2.6
1	A	319	GLU	2.5
1	A	285	LEU	2.5
1	B	196	HIS	2.4
1	A	316	GLU	2.4
1	B	318	ALA	2.4
1	B	301	ASN	2.4
1	B	180	GLN	2.4
1	B	241	THR	2.4
1	A	306	GLN	2.4
1	A	315	GLN	2.4
1	A	241	THR	2.4
1	B	187	ASN	2.3
1	B	188	ASN	2.3
1	B	281	ARG	2.3
1	A	194	GLY	2.3
1	B	339	GLU	2.3
1	B	185	PHE	2.3
1	B	189	THR	2.2
1	B	304	GLU	2.2
1	A	338	ARG	2.2
1	B	302	ALA	2.2
1	B	337	GLU	2.1
1	A	305	SER	2.1
1	A	313	VAL	2.1
1	A	188	ASN	2.1
1	B	223	HIS	2.1
1	B	182	TYR	2.0
1	A	280	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	1SY	A	401[A]	45/45	0.94	0.08	12,15,17,18	45
2	1SY	A	401[B]	45/45	0.94	0.08	12,15,17,18	45

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 1SY A 401 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 1SY A 401 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.