



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:27 AM UTC

PDB ID : 9YZR / pdb\_00009yzt  
Title : Isoreticular co-crystal 1 with asymmetrical expanded duplex (31mer) containing insert sequence TAATTAGGCCG and loaded with ultrabithorax homeodomain  
Authors : Shields, E.T.; Slaughter, C.K.; Magna, E.N.; Snow, C.D.  
Deposited on : 2025-10-30  
Resolution : 3.39 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1001 (3.44-3.36)
Clashscore	190562	1022 (3.44-3.36)
Ramachandran outliers	187476	1012 (3.44-3.36)
Sidechain outliers	187428	1012 (3.44-3.36)
RSRZ outliers	180081	1001 (3.44-3.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	31	<div> <div>42%</div> <div>58%</div> </div>
2	B	31	<div> <div>68%</div> <div>32%</div> </div>
3	C	263	<div> <div>6%</div> <div>73%</div> <div>9%</div> <div>18%</div> </div>
4	D	62	<div> <div>23%</div> <div>73%</div> <div>27%</div> </div>
4	E	62	<div> <div>29%</div> <div>60%</div> <div>40%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	MG	C	301	-	-	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3409 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (31-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	31	Total	C	N	O	P	0	0	0
			631	301	113	186	31			

- Molecule 2 is a DNA chain called DNA (31-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	31	Total	C	N	O	P	0	0	0
			640	304	119	186	31			

- Molecule 3 is a protein called Replication initiation protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	216	Total	C	N	O	S	0	2	0
			1705	1091	294	313	7			

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-11	MET	-	initiating methionine	UNP P03856
C	-10	ARG	-	expression tag	UNP P03856
C	-9	GLY	-	expression tag	UNP P03856
C	-8	SER	-	expression tag	UNP P03856
C	-7	HIS	-	expression tag	UNP P03856
C	-6	HIS	-	expression tag	UNP P03856
C	-5	HIS	-	expression tag	UNP P03856
C	-4	HIS	-	expression tag	UNP P03856
C	-3	HIS	-	expression tag	UNP P03856
C	-2	HIS	-	expression tag	UNP P03856
C	-1	GLY	-	expression tag	UNP P03856
C	0	SER	-	expression tag	UNP P03856
C	118	PRO	ARG	conflict	UNP P03856

- Molecule 4 is a protein called Homeotic protein ultrabithorax.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	45	Total	C	N	O	0	0	0
			244	146	47	51			
4	E	37	Total	C	N	O	0	0	0
			188	112	38	38			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	1	GLY	-	expression tag	UNP P83949
D	2	THR	-	expression tag	UNP P83949
E	1	GLY	-	expression tag	UNP P83949
E	2	THR	-	expression tag	UNP P83949

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	1	Total	Mg	0	0
			1	1		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA (31-MER)

Chain A: 




- Molecule 2: DNA (31-MER)

Chain B: 





- Molecule 3: Replication initiation protein

Chain C: 




- Molecule 4: Homeotic protein ultrabithorax

Chain D: 



- Molecule 4: Homeotic protein ultrabithorax

Chain E: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	76.03Å 124.86Å 139.09Å 90.00° 91.85° 90.00°	Depositor
Resolution (Å)	45.29 – 3.39 45.29 – 3.39	Depositor EDS
% Data completeness (in resolution range)	98.3 (45.29-3.39) 98.3 (45.29-3.39)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.61 (at 3.40Å)	Xtriage
Refinement program	PHENIX 1.21.2_5419	Depositor
R, $R_{free}$	0.265 , 0.298 0.265 , 0.295	Depositor DCC
$R_{free}$ test set	1774 reflections (9.81%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	111.2	Xtriage
Anisotropy	0.150	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 103.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.42$ , $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	0.046 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3409	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	146.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.19% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.16	0/706	0.38	0/1086
2	B	0.19	0/718	0.38	0/1107
3	C	0.12	0/1752	0.40	0/2374
4	D	0.08	0/241	0.21	0/330
4	E	0.07	0/181	0.15	0/242
All	All	0.14	0/3598	0.37	0/5139

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	631	0	350	11	0
2	B	640	0	350	6	0
3	C	1705	0	1586	12	0
4	D	244	0	125	0	0
4	E	188	0	74	0	0
5	C	1	0	0	0	0
All	All	3409	0	2485	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 5.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:20:DA:H2''	1:A:21:DT:H5''	1.67	0.75
2:B:13:DC:H2''	2:B:14:DG:C8	2.45	0.51
1:A:26:DA:H2''	1:A:27:DG:C8	2.46	0.51
1:A:25:DT:H2''	1:A:26:DA:C8	2.48	0.49
2:B:30:DC:H2''	2:B:31:DA:C8	2.47	0.49
2:B:18:DT:H2''	2:B:19:DA:C8	2.49	0.48
1:A:10:DA:H4'	1:A:11:DA:OP1	2.14	0.48
3:C:180:ILE:HA	3:C:240:PHE:O	2.14	0.47
1:A:27:DG:H2''	1:A:28:DG:N7	2.29	0.47
3:C:201:MET:N	3:C:202:PRO:HD2	2.30	0.47
1:A:9:DA:H2''	1:A:10:DA:C8	2.50	0.46
1:A:13:DT:H2'	1:A:14:DG:C8	2.51	0.46
3:C:195:PRO:HD2	3:C:198:TYR:CD2	2.51	0.46
3:C:156:GLU:O	3:C:219:ARG:HD2	2.17	0.45
1:A:1:DC:H2'	1:A:2:DC:C6	2.52	0.44
3:C:57:ASP:CG	3:C:58:GLY:H	2.25	0.44
3:C:15:SER:OG	3:C:151:LEU:HB3	2.18	0.43
3:C:38:MET:HE1	3:C:82:ILE:HA	1.99	0.43
1:A:3:DT:H2'	1:A:4:DG:C8	2.53	0.43
3:C:94:VAL:HB	3:C:114:TRP:CZ2	2.54	0.43
3:C:185:ILE:HG21	3:C:201:MET:HE1	2.01	0.43
1:A:28:DG:H1'	1:A:29:DC:H5'	2.01	0.43
1:A:30:DC:H2''	1:A:31:DG:C8	2.54	0.42
2:B:15:DG:H2''	2:B:16:DC:H5'	2.00	0.42
2:B:26:DA:H2''	2:B:27:DG:H5''	2.02	0.42
3:C:93:GLU:HG2	3:C:113:PRO:HA	2.02	0.42
3:C:198:TYR:O	3:C:204:PHE:HB2	2.21	0.41
2:B:30:DC:H2''	2:B:31:DA:N7	2.35	0.41
3:C:28:ALA:HA	3:C:191:ARG:O	2.21	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	C	210/263 (80%)	205 (98%)	4 (2%)	1 (0%)	24	54
4	D	37/62 (60%)	37 (100%)	0	0	100	100
4	E	23/62 (37%)	20 (87%)	3 (13%)	0	100	100
All	All	270/387 (70%)	262 (97%)	7 (3%)	1 (0%)	30	59

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	143	GLN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	C	174/236 (74%)	174 (100%)	0	100	100
4	D	6/58 (10%)	6 (100%)	0	100	100
4	E	1/58 (2%)	1 (100%)	0	100	100
All	All	181/352 (51%)	181 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	A	31/31 (100%)	-0.23	0	100	100	99, 148, 206, 211	0
2	B	31/31 (100%)	-0.21	0	100	100	93, 143, 208, 211	0
3	C	216/263 (82%)	0.59	15 (6%)	23	18	58, 118, 160, 207	2 (0%)
4	D	45/62 (72%)	1.51	14 (31%)	1	2	175, 236, 280, 317	0
4	E	37/62 (59%)	2.35	18 (48%)	0	1	148, 241, 321, 383	0
All	All	360/449 (80%)	0.75	47 (13%)	7	8	58, 129, 266, 383	2 (0%)

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	E	46	GLN	9.0
4	E	47	ILE	8.7
4	D	56	MET	8.1
4	E	45	ARG	7.3
4	D	54	ARG	6.2
4	E	43	THR	5.5
4	D	55	ARG	5.2
4	E	7	ARG	5.1
4	D	53	ASN	5.0
4	D	47	ILE	4.9
4	D	48	LYS	4.5
3	C	67	TYR	4.4
4	D	50	TRP	4.2
4	E	14	GLN	4.0
4	E	10	TYR	3.4
4	D	52	GLN	3.4
4	D	58	LEU	3.2
3	C	37	ARG	3.1
4	D	41	CYS	3.1
4	E	42	LEU	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
4	D	7	ARG	3.0
4	D	51	PHE	3.0
3	C	233	ARG	2.9
4	E	21	GLU	2.8
3	C	116	ILE	2.8
3	C	49	SER	2.8
3	C	211	VAL	2.8
4	E	41	CYS	2.8
3	C	215	GLU	2.8
4	D	10	TYR	2.7
3	C	84	GLN	2.7
3	C	61[A]	GLU	2.7
3	C	63	HIS	2.7
4	E	49	ILE	2.6
3	C	40	TYR	2.6
4	E	13	TYR	2.6
3	C	96	PHE	2.6
4	E	53	ASN	2.6
4	E	35	GLU	2.6
4	E	15	THR	2.6
3	C	14	ASN	2.5
4	E	50	TRP	2.4
3	C	135	LEU	2.3
4	E	44	GLU	2.3
3	C	139	PHE	2.2
4	E	38	HIS	2.1
4	D	45	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	MG	C	301	1/1	0.62	0.42	72,72,72,72	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.