



## Full wwPDB EM Validation Report ⓘ

Apr 5, 2026 – 11:28 PM UTC

PDB ID : 9ZQB / pdb\_00009zqb  
EMDB ID : EMD-74563  
Title : Nucleosome with an SSB at SHL -2.8 in complex with human PARP2 and HPF1, Class 1  
Authors : Kim, T.H.; Jayathilake, C.; Virk, R.K.; Gregory-Lott, E.R.  
Deposited on : 2025-12-18  
Resolution : 2.10 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132  
MolProbity : 4-5-2 with Phenix2.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
EM percentile statistics : 202505.v01 (Using data in the EMDb archive up until May 2025)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49



# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.













Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	2317 ( 1.60 - 2.60 )

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	124	85% 13%
1	B	124	85% 12%
2	C	125	76% 22%
2	D	125	74% 22%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	E	136	
3	F	136	
4	G	104	
4	H	104	
5	I	274	
5	J	274	
6	K	69	
7	L	128	
8	M	197	
9	P	494	

## 2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 31635 atoms, of which 14529 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone H2A.

Mol	Chain	Residues	Atoms						AltConf	Trace
1	A	108	Total	C	H	N	O	S	0	0
			1708	520	883	161	143	1		
1	B	109	Total	C	H	N	O	S	0	0
			1730	526	896	163	144	1		

- Molecule 2 is a protein called Histone H2B.

Mol	Chain	Residues	Atoms						AltConf	Trace
2	C	98	Total	C	H	N	O	S	0	0
			1605	492	825	143	143	2		
2	D	98	Total	C	H	N	O	S	0	0
			1605	492	825	143	143	2		

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	GLY	-	expression tag	UNP P02283
C	2	ILE	-	insertion	UNP P02283
D	0	GLY	-	expression tag	UNP P02283
D	2	ILE	-	insertion	UNP P02283

- Molecule 3 is a protein called Histone H3.

Mol	Chain	Residues	Atoms						AltConf	Trace
3	E	100	Total	C	H	N	O	S	0	0
			1695	520	870	160	143	2		
3	F	99	Total	C	H	N	O	S	0	0
			1673	514	857	158	142	2		

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	111	SER	CYS	conflict	UNP P02299

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
F	111	SER	CYS	conflict	UNP P02299

- Molecule 4 is a protein called Histone H4.

Mol	Chain	Residues	Atoms						AltConf	Trace
4	G	86	Total	C	H	N	O	S	0	0
			1437	436	743	140	117	1		
4	H	86	Total	C	H	N	O	S	0	0
			1437	436	743	140	117	1		

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	2	ILE	-	insertion	UNP P84040
H	2	ILE	-	insertion	UNP P84040

- Molecule 5 is a protein called mAb PL2-6 antibody.

Mol	Chain	Residues	Atoms						AltConf	Trace
5	I	231	Total	C	H	N	O	S	0	0
			3559	1137	1753	297	362	10		
5	J	231	Total	C	H	N	O	S	0	0
			3559	1137	1753	297	362	10		

- Molecule 6 is a DNA chain called DNA (69-MER).

Mol	Chain	Residues	Atoms						AltConf	Trace
6	K	47	Total	C	H	N	O	P	0	0
			1489	455	529	172	286	47		

- Molecule 7 is a DNA chain called DNA (128-MER).

Mol	Chain	Residues	Atoms						AltConf	Trace
7	L	106	Total	C	H	N	O	P	0	0
			3345	1025	1191	385	638	106		

- Molecule 8 is a DNA chain called DNA (197-MER).

Mol	Chain	Residues	Atoms						AltConf	Trace
8	M	153	Total	C	H	N	O	P	0	0
			4874	1494	1714	600	913	153		


- Molecule 9 is a protein called Poly [ADP-ribose] polymerase 2.


Mol	Chain	Residues	Atoms						AltConf	Trace
9	P	118	Total	C	H	N	O	S	0	0
			1919	615	947	169	180	8		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Histone H2A

Chain A:  85% 13%





- Molecule 1: Histone H2A

Chain B:  85% 12%




- Molecule 2: Histone H2B

Chain C:  76% 22%




- Molecule 2: Histone H2B

Chain D:  74% 22%



- Molecule 3: Histone H3

Chain E:  72% 26%



- Molecule 3: Histone H3





DG	C1
DA	T2
DC	C3
DC	T4
DA	
DC	T24
DC	A25
DC	C26
DC	G27
DC	C28
DC	C29
DC	C30
DC	T31
	C34
	C35
	T45
	A46
	G50
	C51
	T60
	A62
	C63
	T64
	C67
	T68
	A69
	C70
	T71
	C79
	A80
	G84
	T85
	T101
	G102
	T103
	G104
	A105
	T106
	DT
	DG
	DT
	DA
	DT
	DT
	DG
	DA
	DA
	DC
	DG
	DC

[illegible][illegible]

## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	952082	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	49	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	1800	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.633	Depositor
Minimum map value	-0.176	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.012	Depositor
Recommended contour level	0.0369	Depositor
Map size (Å)	331.2, 331.2, 331.2	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.828, 0.828, 0.828	Depositor

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.20	0/835	0.30	0/1124
1	B	0.21	0/844	0.34	0/1135
2	C	0.22	0/791	0.32	0/1057
2	D	0.21	0/791	0.29	0/1057
3	E	0.20	0/837	0.33	0/1120
3	F	0.19	0/828	0.30	0/1109
4	G	0.20	0/702	0.33	0/937
4	H	0.20	0/702	0.31	0/937
5	I	0.20	0/1848	0.40	0/2502
5	J	0.20	0/1848	0.36	0/2502
6	K	0.25	0/1074	0.46	0/1655
7	L	0.27	0/2411	0.47	0/3712
8	M	0.26	0/3551	0.43	0/5484
9	P	0.18	0/993	0.37	0/1334
All	All	0.23	0/18055	0.39	0/25665

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	825	883	882	1	0
1	B	834	896	895	5	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	780	825	824	1	0
2	D	780	825	824	5	0
3	E	825	870	869	1	0
3	F	816	857	856	2	0
4	G	694	743	742	3	0
4	H	694	743	742	1	0
5	I	1806	1753	1751	19	0
5	J	1806	1753	1751	11	0
6	K	960	529	529	5	0
7	L	2154	1191	1192	17	0
8	M	3160	1714	1715	11	0
9	P	972	947	946	7	0
All	All	17106	14529	14518	86	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (86) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:I:24:MET:HA	5:I:24:MET:HE2	1.52	0.91
5:I:132:ARG:NH1	5:I:132:ARG:HA	2.10	0.67
4:G:26:ASP:OD1	4:G:26:ASP:O	2.18	0.62
1:B:13:LYS:HA	1:B:13:LYS:HE2	1.85	0.59
5:J:174:MET:HE1	5:J:248:TYR:HB2	1.86	0.58
9:P:157:MET:N	9:P:157:MET:HE2	2.19	0.56
5:I:167:GLU:N	5:I:167:GLU:OE1	2.39	0.56
5:I:196:ARG:HB3	5:I:196:ARG:NH1	2.21	0.55
5:I:216:GLU:HA	5:I:216:GLU:OE2	2.07	0.53
8:M:52:DC:H2''	8:M:53:DT:H72	1.91	0.53
5:J:155:GLU:N	5:J:155:GLU:OE2	2.42	0.52
5:J:167:GLU:N	5:J:167:GLU:OE1	2.42	0.52
8:M:148:DC:H2''	8:M:149:DA:C8	2.45	0.52
5:J:243:GLU:OE2	5:J:243:GLU:C	2.55	0.50
6:K:34:DT:H2'	6:K:35:DT:H71	1.93	0.50
5:J:196:ARG:HB2	5:J:196:ARG:NH1	2.26	0.50
5:J:129:GLU:N	5:J:129:GLU:OE1	2.44	0.50
7:L:101:DT:H2''	7:L:102:DG:C8	2.46	0.50
9:P:195:GLU:N	9:P:195:GLU:OE2	2.45	0.50
5:I:174:MET:HE1	5:I:248:TYR:HB2	1.95	0.49
5:J:28:MET:HE3	5:J:47:CYS:SG	2.52	0.49

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:L:50:DG:H2''	7:L:51:DC:O5'	2.11	0.49
8:M:97:DG:H2''	8:M:98:DG:O5'	2.12	0.49
5:I:166:VAL:HG11	5:I:240:LEU:HD13	1.95	0.49
8:M:53:DT:H2'	8:M:54:DT:H71	1.94	0.49
8:M:84:DC:H2''	8:M:85:DG:C8	2.48	0.49
7:L:26:DC:H2''	7:L:27:DG:C8	2.48	0.48
7:L:60:DT:H2'	7:L:61:DT:H71	1.95	0.48
6:K:34:DT:C2'	6:K:35:DT:H71	2.44	0.47
5:J:217:LYS:NZ	5:J:217:LYS:HB3	2.29	0.47
8:M:97:DG:H4'	8:M:98:DG:OP1	2.15	0.47
9:P:133:LEU:C	9:P:133:LEU:HD23	2.40	0.47
2:D:79:LEU:HD11	2:D:95:THR:HG22	1.97	0.46
1:B:54:VAL:HG21	2:D:97:VAL:HG21	1.97	0.46
5:I:154:MET:C	5:I:154:MET:SD	2.99	0.46
5:I:24:MET:HA	5:I:24:MET:CE	2.36	0.46
7:L:50:DG:H4'	7:L:51:DC:OP1	2.16	0.46
5:I:80:ASP:O	5:I:80:ASP:OD2	2.34	0.46
7:L:63:DC:H2'	7:L:64:DT:H72	1.99	0.45
7:L:103:DT:H2''	7:L:104:DG:C8	2.52	0.45
8:M:23:DC:H2''	8:M:24:DA:C8	2.51	0.45
5:I:196:ARG:HB3	5:I:196:ARG:CZ	2.46	0.45
1:B:91:GLU:OE1	5:I:208:TYR:CD1	2.69	0.45
4:H:25:ARG:NH1	4:H:25:ARG:HB2	2.31	0.45
6:K:29:DC:H2''	6:K:30:DT:H71	1.99	0.45
2:D:79:LEU:HD11	2:D:95:THR:CG2	2.47	0.45
7:L:34:DC:H2''	7:L:35:DC:C5	2.51	0.45
8:M:5:DC:H2''	8:M:6:DA:C8	2.51	0.45
3:E:66:LEU:HB3	3:E:67:PRO:HD3	1.99	0.45
5:I:72:ILE:HG22	5:I:74:TYR:O	2.16	0.45
5:I:154:MET:O	5:I:155:GLU:OE2	2.35	0.45
9:P:174:GLU:HA	9:P:174:GLU:OE1	2.16	0.45
5:I:176:CYS:SG	5:I:250:CYS:SG	3.11	0.44
7:L:30:DC:H2'	7:L:31:DT:H71	1.99	0.44
7:L:79:DC:H2''	7:L:80:DA:C8	2.53	0.44
7:L:45:DT:H2''	7:L:46:DA:C8	2.53	0.44
3:F:91:MET:HA	3:F:91:MET:HE2	1.98	0.44
5:I:129:GLU:OE1	5:I:129:GLU:C	2.60	0.44
8:M:68:DC:H2''	8:M:69:DG:C8	2.52	0.44
3:F:38:LYS:N	3:F:39:PRO:CD	2.82	0.43
7:L:3:DC:H2''	7:L:4:DT:H71	2.01	0.43
8:M:131:DT:H2''	8:M:132:DC:C5	2.54	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:30:ARG:HE	2:D:30:ARG:H	1.67	0.42
1:B:61:GLU:OE2	1:B:61:GLU:HA	2.20	0.42
2:C:85:ARG:HG3	2:C:85:ARG:HH11	1.84	0.42
7:L:67:DC:C2'	7:L:68:DT:H72	2.49	0.42
6:K:20:DC:H2''	6:K:21:DC:C6	2.54	0.42
7:L:84:DG:H2''	7:L:85:DT:C6	2.54	0.42
7:L:24:DT:H2''	7:L:25:DA:N7	2.35	0.42
7:L:28:DC:H2''	7:L:29:DG:C8	2.55	0.42
7:L:70:DC:H2'	7:L:71:DT:H71	2.00	0.42
9:P:110:GLU:CD	9:P:110:GLU:O	2.63	0.42
4:G:25:ARG:HD2	4:G:26:ASP:HB3	2.01	0.42
5:J:65:TRP:C	5:J:66:LYS:HD2	2.45	0.42
6:K:29:DC:H2''	6:K:30:DT:C7	2.50	0.41
1:A:76:THR:O	1:A:76:THR:HG22	2.21	0.41
1:B:61:GLU:OE2	5:I:257:LEU:HD12	2.21	0.41
2:D:30:ARG:O	2:D:30:ARG:HD2	2.21	0.41
5:I:74:TYR:O	5:I:75:ALA:C	2.62	0.41
5:J:192:LYS:HB2	5:J:202:ILE:HD11	2.02	0.41
9:P:202:ASP:OD1	9:P:203:MET:N	2.53	0.41
4:G:26:ASP:O	4:G:26:ASP:CG	2.63	0.41
5:I:131:LYS:N	5:I:131:LYS:HD2	2.36	0.41
5:J:66:LYS:HA	5:J:66:LYS:HE2	2.03	0.40
8:M:92:DT:H2''	8:M:93:DA:C8	2.57	0.40
9:P:124:LEU:HD22	9:P:124:LEU:N	2.37	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	106/124 (86%)	105 (99%)	1 (1%)	0	100	100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	107/124 (86%)	105 (98%)	2 (2%)	0	100	100
2	C	96/125 (77%)	96 (100%)	0	0	100	100
2	D	96/125 (77%)	96 (100%)	0	0	100	100
3	E	98/136 (72%)	97 (99%)	1 (1%)	0	100	100
3	F	97/136 (71%)	97 (100%)	0	0	100	100
4	G	84/104 (81%)	82 (98%)	2 (2%)	0	100	100
4	H	84/104 (81%)	82 (98%)	2 (2%)	0	100	100
5	I	227/274 (83%)	224 (99%)	3 (1%)	0	100	100
5	J	227/274 (83%)	221 (97%)	6 (3%)	0	100	100
9	P	116/494 (24%)	113 (97%)	3 (3%)	0	100	100
All	All	1338/2020 (66%)	1318 (98%)	20 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	83/94 (88%)	81 (98%)	2 (2%)	43	49
1	B	84/94 (89%)	84 (100%)	0	100	100
2	C	85/105 (81%)	83 (98%)	2 (2%)	43	49
2	D	85/105 (81%)	84 (99%)	1 (1%)	63	72
3	E	87/111 (78%)	87 (100%)	0	100	100
3	F	86/111 (78%)	85 (99%)	1 (1%)	63	72
4	G	71/80 (89%)	71 (100%)	0	100	100
4	H	71/80 (89%)	71 (100%)	0	100	100
5	I	202/226 (89%)	202 (100%)	0	100	100
5	J	202/226 (89%)	202 (100%)	0	100	100
9	P	105/433 (24%)	104 (99%)	1 (1%)	68	76

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1161/1665 (70%)	1154 (99%)	7 (1%)	76 86

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	41	GLU
1	A	101	THR
2	C	47	VAL
2	C	100	LEU
2	D	56	LYS
3	F	98	GLU
9	P	116	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	24	GLN
1	A	68	ASN
1	A	110	ASN
1	A	112	GLN
1	B	24	GLN
1	B	104	GLN
1	B	112	GLN
2	C	48	HIS
2	D	66	ASN
2	D	94	GLN
5	I	159	GLN
5	J	30	GLN
5	J	115	HIS
9	P	121	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.



## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-74563. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 200




Y Index: 200




Z Index: 200


### 6.2.2 Raw map



X Index: 200



Y Index: 200




Z Index: 200


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 199




Y Index: 183




Z Index: 219


### 6.3.2 Raw map



X Index: 199



Y Index: 183




Z Index: 219


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) ⓘ


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0369. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

## 7.2 Volume estimate [i](#)




The volume at the recommended contour level is 230 nm<sup>3</sup>; this corresponds to an approximate mass of 208 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



### 7.3 Rotationally averaged power spectrum ⓘ




\*Reported resolution corresponds to spatial frequency of 0.476 Å<sup>-1</sup>

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.476  $\text{\AA}^{-1}$

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.10	-	-
Author-provided FSC curve	2.08	2.37	2.12
Unmasked-calculated*	2.67	3.03	2.72

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 2.67 differs from the reported value 2.1 by more than 10 %


## 9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-74563 and PDB model 9ZQB. Per-residue inclusion information can be found in section [3](#) on page [7](#).


### 9.1 Map-model overlay [i](#)



X




Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.0369 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0369).































## 9.4 Atom inclusion [i](#)



At the recommended contour level, 99% of all backbone atoms, 99% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.0369) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9900	 0.6190
A	 0.9860	 0.6590
B	 0.9880	 0.6540
C	 0.9720	 0.6440
D	 0.9870	 0.6450
E	 0.9790	 0.6530
F	 0.9820	 0.6460
G	 0.9530	 0.6420
H	 0.9820	 0.6470
I	 0.9930	 0.5960
J	 0.9940	 0.6060
K	 1.0000	 0.6070
L	 1.0000	 0.6190
M	 1.0000	 0.6140
P	 0.9870	 0.5130

